




Chameleon Drive, Erskine Park
Biodiversity Development Assessment Report

Penrith City Council

DOCUMENT TRACKING

Project Name	Biodiversity Development Assessment Report: Chameleon Drive, Erskine Park
Project Number	21SUT-20205
Project Manager	Karen Spicer
Accredited Assessor Certification	 Karen Spicer BAAS18141
Prepared by	Annabelle McTaggart and Karen Spicer
Reviewed by	Michelle Frolich
Approved by	Deanne Hickey
Status	Final
Version Number	V2
Last saved on	31 January 2022

This report should be cited as 'Eco Logical Australia 2022. *Biodiversity Development Assessment Report: Chameleon Drive, Erskine Park.* Prepared for Penrith City Council.'

ACKNOWLEDGEMENTS

This document has been prepared by Eco Logical Australia Pty Ltd with support from Karen Luka and Vince Hardy

Disclaimer

This document may only be used for the purpose for which it was commissioned and in accordance with the contract between Eco Logical Australia Pty Ltd and Penrith City Council. The scope of services was defined in consultation with Penrith City Council, by time and budgetary constraints imposed by the client, and the availability of reports and other data on the subject area. Changes to available information, legislation and schedules are made on an ongoing basis and readers should obtain up to date information. Eco Logical Australia Pty Ltd accepts no liability or responsibility whatsoever for or in respect of any use of or reliance upon this report and its supporting material by any third party. Information provided is not intended to be a substitute for site specific assessment or legal advice in relation to any matter. Unauthorised use of this report in any form is prohibited.

Template 2.B.1

Executive Summary

Eco Logical Australia Pty Ltd (ELA) was engaged by Penrith City Council to prepare a Streamlined (small area) Biodiversity Development Assessment Report (BDAR) for Lot 1106 DP709078 and Lot 1107 DP709078, Chameleon Dr, Erskine Park (the development site) within the Penrith local government area (LGA).

This report has been prepared consistent with the Biodiversity Assessment Method (BAM) 2020, under the NSW Biodiversity Conservation Act 2016 (BC Act).

The development site (Lot 1106 DP709078 and part Lot 1107 DP709078) is 0.56 ha. The proposed development will facilitate a 5-lot subdivision within part of Chameleon Reserve. Prior to site development, the clearance of all vegetation within the development footprint will be required to allow for the construction of the four residences and associated driveways and landscaped areas. Impacts to trees adjacent to the development site have been assessed by Laws 2021 and tree retention/removal has been excluded/included in the development footprint based on this assessment.

Field survey was undertaken to validate and map vegetation, undertake the required vegetation integrity plots and assess habitat values. The development footprint is 0.27 ha and consists of 0.003 ha of planted native vegetation, 0.01 ha of exotic vegetation and 0.03 ha of PCT 849 *Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion*. PCT 849 was present in one condition state – low. The remainder of the development footprint (0.23 ha) is cleared land.

The 0.003 ha of planted native vegetation was assessed against Appendix D: *Streamlined assessment module - planted native vegetation* of the BAM 2020 and does not require offsetting. The remaining vegetation within the development site was either exotic to NSW or cleared land dominated by exotic grasses.

One full-floristic vegetation integrity plot was undertaken within PCT 849_low in accordance with the BAM. A vegetation integrity assessment using the BAM Calculator (BAMC) was undertaken and resulted in a vegetation integrity score of 9.4, which is too low to require offsets.

PCT 849 forms a part of the threatened ecological community (TEC) *Cumberland Plain Woodland in the Sydney Basin Bioregion*. This vegetation community is listed as Critically Endangered under the BC Act. PCT 849 also forms a part of the Federally listed TEC *Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest*. However, the vegetation within the development site did not meet any of the required condition classes and as such, is not recognised under the EPBC Act.

Cumberland Plain Woodland in the Sydney Basin Bioregion is a candidate for a Serious and Irreversible Impact (SAIL) and an assessment has been undertaken to assist the consent authority to decide whether any SAIL will result.

As this BDAR is being assessed via the streamlined (small area) assessment (BAM 2020, Appendix C), only those candidate species at risk of a serious and irreversible impact (SAIL) require targeted survey. All the candidate species generated by the BAMC are entities at risk of a SAIL. However, the development site does not contain breeding habitat for the candidate fauna species (*Anthochaera Phrygia*, *Chalinolobus dwyeri*, *Lathamus discolor*, *Miniopterus australis*, *Miniopterus orianae oceanensis*). Only

one flora species, *Caladenia tessellata*, was considered a candidate species. However, the development site does not provide habitat for this species. As such, no further assessment or targeted survey is required, and no species credits are required to offset impacts.

Supervision of vegetation removal by an ecologist should be undertaken to ensure any fauna present are taken into care and relocated into similar nearby habitat. Sediment and erosion control measures are recommended to prevent stormwater pollution or movement of soils from the development site.

Contents

- 1. Introduction 1**
- 1.1. General description of the development site 1
- 1.2. Brief description of the proposal 1
- 1.3. Development site footprint 1
- 1.4. Sources of information used 2
- 1.5. Legislative context 5

- 2. Landscape features 7**
- 3. Native Vegetation 8**
- 3.1. Survey Effort 8
- 3.2. Native vegetation extent on the development site 8
- 3.3. Plant Community Types present 10
- 3.3.1. Plant Community Type selection justification 10
- 3.4. Threatened Ecological Communities 11
- 3.5. Vegetation integrity assessment 11
- 3.5.1. Vegetation zones 11
- 3.5.2. Decision-making key for planted native vegetation (Appendix D BAM 2020) 13
- 3.5.3. Patch size 15
- 3.5.4. Assessing vegetation integrity 16

- 4. Threatened species 20**
- 4.1. Ecosystem credit species 20
- 4.2. Species credit species 22
- 4.2.1. Identification of species credit species 22
- 4.2.2. Assessment of habitat constraints and vagrant species 24
- 4.2.3. Candidate species requiring further assessment 24
- 4.3. Identification of prescribed additional biodiversity impact entities 25
- 4.3.1. Karst, caves, crevices, cliffs, rocks and other geological features of significance 25
- 4.3.2. Human-made structures and non-native vegetation 25
- 4.3.3. Habitat connectivity 25
- 4.3.4. Water bodies, water quality and hydrological processes 25
- 4.3.5. Wind farm developments 25
- 4.3.6. Vehicle strikes 25

- 5. Avoiding and Minimising Impacts on Biodiversity Values 26**
- 5.1. Locating a project to avoid and minimise impacts on biodiversity values 26
- 5.1.1. Direct and indirect impacts 26
- 5.1.2. Prescribed biodiversity impacts 27
- 5.2. Designing a project to avoid and minimise impacts on biodiversity values 27

5.2.1. Direct and indirect impacts	27
5.2.2. Prescribed biodiversity impacts	27
6. Assessment of Impacts	28
6.1. Direct impacts	28
6.2. Change in vegetation integrity	28
6.3. Indirect impacts	30
6.4. Prescribed biodiversity impacts	31
6.5. Mitigating and managing direct and indirect impacts	33
6.6. Mitigating prescribed impacts	36
6.7. Adaptive management strategy	38
7. Impact summary	39
7.1. Serious and Irreversible Impacts (SAIL)	39
7.2. Impacts requiring offsets	42
7.3. Impacts not requiring offsets	42
7.4. Areas not requiring assessment	43
7.5. Credit summary	43
8. Consistency with legislation and policy	45
8.1. <i>Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act)</i>	45
8.1.1. Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest ecological community	45
8.1.2. <i>Pteropus poliocephalus</i> (Grey-headed Flying-fox)	47
9. Conclusion	49
10. References	50
Appendix A Definitions	52
Appendix B Vegetation Floristic Plot Data	55
Appendix C Vegetation Integrity Plot Data	57
Appendix D Biodiversity credit report	58
Appendix E Likelihood of occurrence	60

List of Figures

Figure 1: Location map and landscape features	3
Figure 2: Site plan.....	4
Figure 3: Survey effort.....	9
Figure 4: Plant Community Types	17
Figure 5: Threatened Ecological Communities.....	18
Figure 6: Vegetation Zones and Plots.....	19
Figure 7: Combined impact map	29
Figure 8: Serious and Irreversible Impacts.....	44
Figure 9: Flowchart to assist with the determination of EPBC Act condition thresholds (Commonwealth of Australia, 2010).....	46

List of Tables

Table 1: Legislative context.....	5
Table 2: Landscape features.....	7
Table 3: Full-floristic PCT identification plots.....	8
Table 4: Plant community types.....	10
Table 5: Potential PCTs.....	10
Table 6: Threatened Ecological Communities within the development footprint	11
Table 7: Vegetation zones and vegetation integrity survey plots collected on the development site....	11
Table 8: Description of the vegetation zones	12
Table 9: Vegetation integrity scores	16
Table 10: Predicted ecosystem credit species	20
Table 11: Justification for exclusion of predicted ecosystem credit species	22
Table 12: Candidate species credit species.....	22
Table 13: Justification for the exclusion/ inclusion of species credit species.....	23
Table 14: Direct impacts to native vegetation	28
Table 15: Change in vegetation integrity	28
Table 16: Indirect impacts.....	30
Table 17: Direct impacts on prescribed biodiversity impacts.....	31
Table 18: Measures proposed to mitigate and manage impacts.....	33
Table 19: Mitigation measures for prescribed biodiversity impacts	36
Table 20: Serious and Irreversible Impacts Summary.....	39
Table 21: Evaluation of an impact on a TEC consistent with 9.1.1 of the BAM	39
Table 22: Impacts to native vegetation that do not require offsets.....	43
Table 23: Ecosystem credits required	43
Table 24: EPBC Act impact assessment for <i>Pteropus poliocephalus</i> (Grey-headed Flying-fox).....	47

Abbreviations

Abbreviation	Description
BAM	Biodiversity Assessment Method
BAMC	Biodiversity Assessment Method Credit Calculator
BC Act	NSW <i>Biodiversity Conservation Act 2016</i>
BDAR	Biodiversity Development Assessment Report
CEEC	Critically Endangered Ecological Community
DAWE	Commonwealth Department of Agriculture, Water and the Environment
DPIE	NSW Department of Planning, Industry and Environment
EEC	Endangered Ecological Community
ELA	Eco Logical Australia Pty Ltd
EP&A Act	NSW <i>Environmental Planning and Assessment Act 1979</i>
EPBC Act	Commonwealth <i>Environment Protection and Biodiversity Conservation Act 1999</i>
FM Act	NSW <i>Fisheries Management Act 1994</i>
GIS	Geographic Information System
GPS	Global Positioning System
IBRA	Interim Biogeographic Regionalisation for Australia
LGA	Local Government Area
LLS	Local Land Service
NSW	New South Wales
PCT	Plant Community Type
OEH	Office of Environment and Heritage
SEPP	State Environmental Planning Policy
TEC	Threatened Ecological Community
VIS	Vegetation Information System
WM Act	NSW <i>Water Management Act 2000</i>

1. Introduction

This Biodiversity Development Assessment Report (BDAR) has been prepared by Karen Spicer, an Accredited Person (BAAS18141) to apply the Biodiversity Assessment Method (BAM) under the NSW *Biodiversity Conservation Act 2016* (BC Act). All credit calculations have been undertaken using the BAM Calculator (BAMC) version in case number 00029977/BAAS18141/21/00029978. Given the minimum lot size for the development site is 550 m² and 0.03 ha of native vegetation will be impacted, *Appendix C Streamlined assessment module – Small area* of the BAM was applied.

Definitions of terminology used throughout this report are presented in Appendix A.

1.1. General description of the development site

The development site is located within Chameleon Reserve (Lot 1106 DP709078 and Lot 1107 DP709078) at Chameleon Drive Erskine Park, within Penrith LGA. The development site is zoned R2 Low Density Residential and RE1 Public Recreation under the Penrith Local Environment Plan 2010. Adjacent land to the north and west within Chameleon Reserve is zoned RE1 Public Recreation and all other surrounding land is zoned R2.

The development site consists of cleared exotic parkland with patches of native vegetation, some of which has been planted. Existing residential areas occur to the east, Chameleon Drive forms the southern boundary and Chameleon Reserve continues to the north and west. Chameleon Reserve includes native vegetation, planted native vegetation, a carpark and playing courts.

This report includes two base maps, the Location Map (Figure 1) and the Site Plan (Figure 2).

1.2. Brief description of the proposal

The development site (Lot 1106 DP709078 and Lot 1107 DP709078) is approximately 0.56 ha and is proposed to be subdivided into five lots to facilitate the development of four house and land packages within the south eastern part of the reserve. The land proposed for residential development (referred to as the “development footprint”) has already been rezoned from RE1 to R2 and has a total area of 0.27 ha. The development will include clearing of vegetation to allow for the construction of the new residences, provision of services (water, sewer and electricity), driveways and landscaping.

The project is part of Penrith City Council’s Open Space Reinvestment project. Funds raised from the sale of these packages will be reinvested into open space improvements within the suburb of Erskine Park.

1.3. Development site footprint

The four residential lots will be 562 m², 566 m², 693 m² and 680 m² in area, and will require removal of all vegetation from within the lots (the Development Site Footprint) to allow for construction of the residences, driveways and landscaped areas.

Impacts associated with the proposed development have been assessed for all trees within and immediately adjacent to the development site by Laws 2021. The findings in this Arboricultural Impact Assessment Report (Laws 2021) have been used to determine the extent of impacts for BAMC credit

calculations and preparation of this BDAR. As such, canopy of adjacent vegetation not impacted by the proposed development that overhangs the development footprint has not been included in the impact area/development footprint.

The development site boundary and development footprint are presented in Figure 2.

1.4. Sources of information used

The following data sources were reviewed as part of this report:

- BioNet Vegetation Classification (DPIE 2021)
- Bionet Atlas Database (DPIE 2021) (accessed 29 September 2021)
- Threatened Biodiversity Data Collection
- ePlanning Spatial Viewer (DPIE 2021) (accessed 6 December 2021)
- Protected Matters Search Tool (PMST) (Department of the Environment, Water, Heritage and the Arts (DAWE) 2021)
- National Flying-fox Monitoring Viewer (DAWE 2021) (accessed 6 December 2021)
- Biodiversity Values Map (DPIE 2021) (accessed 6 December 2021)
- *Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest, A guide to identifying and protecting the nationally threatened ecological community Environment Protection and Biodiversity Conservation Act 1999, Policy Statement 3.31.* (DAWE 2010)
- Cumberland Plain Recovery Plan (Department of Environment, Climate Change and Water NSW (DECCW) 2010.
- Remnant Vegetation of the western Cumberland subregion, OEH 2013 Update. VIS_ID 4207




Figure 1: Location map and landscape features




Figure 2: Site plan

1.5. Legislative context

Legislation relevant to the development site is outlined in Table 1.

Table 1: Legislative context

Name	Relevance to the project	Report Section
Commonwealth		
<i>Environmental Protection and Biodiversity Conservation Act 1999</i>	<p>Matters of National Environmental Significance (MNES) have been identified on or near the development site. This report assesses impacts to MNES and concludes that the development is not likely to have a significant impact on MNES. There are seven MNES that are triggers for Commonwealth assessment and approval. These are:</p> <ol style="list-style-type: none"> 1. World Heritage properties 2. National Heritage places 3. Ramsar wetlands of international importance 4. Nationally threatened species and communities 5. Migratory species 6. Nuclear actions 7. Commonwealth marine environment. <p>Appendix E contains a likelihood or occurrence table for species listed under the EPBC Act. One MNES has been identified as having the potential to occur or is known to occur within the development site, <i>Pteropus poliocephalus</i> (Grey-headed Flying-fox). Further assessment of this MNES is addressed in Section 8.1. Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest is not present at the site.</p>	8.1
State		
<i>Environmental Planning and Assessment Act 1979</i>	<p>The EP&A Act is the principal planning legislation for NSW. It provides a framework for the overall environmental planning and assessment of development proposals.</p> <p>The proposed development requires consent under the (Penrith Local Environmental Plan (LEP) and is to be assessed under Part 4 of the EP&A Act.</p>	
<i>Biodiversity Conservation Act 2016</i>	<p>The proposed development will affect land mapped under the Biodiversity Values Map (BV Map) (Figure 2) and therefore requires submission of a Biodiversity Development Assessment Report (BDAR). This BDAR has been prepared consistent with the Biodiversity Assessment Method (BAM) 2020.</p> <p>One entity at risk of a Serious and Irreversible (SAIL) has been identified within the development site - <i>Cumberland Plain Woodland in the Sydney Basin Bioregion</i>. An assessment is provided for assisting council with deciding whether a SAIL is likely to occur</p>	Section 1 to 7
<i>Local Land Services Amendment Act 2016</i>	The Local Land Services Amendment Act 2016 does not apply to areas of the state to which the Vegetation SEPP applies. The Vegetation SEPP applies to Penrith Council local government area.	N/A
<i>Fisheries Management Act 1994</i>	The development does not involve impacts to Key Fish Habitat and does not involve harm to marine vegetation, dredging, reclamation or obstruction of fish passage. A permit or consultation under the FM Act is not required.	N/A
<i>Water Management Act 2000</i>	A controlled activity approval is required under the Water Management Act (WM Act) for some types of activities on waterfront land. Waterfront land is defined as <i>the bed of any river, lake or estuary, and the land within 40 metres of the river banks, lake shore or estuary mean high water mark</i> .	N/A

Name	Relevance to the project	Report Section
	<p>A first order water course occurs to the north of the development site, within 40 m of the development footprint. There are no formed banks or bed and the drainage line is maintained as a mown drainage swale. As such, this water course would not meet the definition of a “river” under the WM Act.</p> <p>In addition, Clause 41 of the <i>Water Management (General) Regulation 2018</i> provides a controlled activity approval exemption to public authorities including local councils. This exemption applies to any kind of controlled activity.</p> <p>As such, a Controlled Activity Approval under s91 of the WM Act is not required.</p>	

Planning Instruments

State Environmental Planning Policy (Coastal Management) 2018	The proposed development is not located on or adjacent to land subject to this SEPP therefore this SEPP is not applicable.	N/A
State Environmental Planning Policy (Koala Habitat Protection)	The State Environmental Planning Policy (Koala Habitat Protection) 2020 and 2021 do not apply to Penrith Local Government Area.	N/A
Penrith Local Environment Plan (LEP) 2010	<p>The development site is zoned R2 Low Density Residential under the Penrith LEP.</p> <p>The development site is not subject to a Biodiversity or Riparian overlay under the LEP.</p>	
Penrith Development Control Plan (DCP) 2014	The Penrith DCP contains provisions relating to native vegetation. Part C2 Vegetation Management sets objectives which includes protection and conservation of biodiversity values of trees and other vegetation in the City. There are no additional requirements under this DCP that must be addressed as part of this BDAR.	

2. Landscape features

The site-based method was applied for this assessment, therefore the assessment area is the 1,500 m buffer surrounding the outside edge of the boundary of the development site. The landscape features considered for this assessment are presented in Table 2, Figure 1 and Figure 2.

Table 2: Landscape features

Landscape feature	Description	Data source
IBRA Region(s)	Sydney Basin	Interim Biogeographic Regionalisation for Australia, Version 7
IBRA subregion(s)	Cumberland	Interim Biogeographic Regionalisation for Australia, Version 7
Rivers and streams	There are no mapped water courses within the development site.	NSW LPI Waterway mapping
Estuaries and wetlands	None present within the development site.	NSW directory of important wetlands
Connectivity of different areas of habitat	The development site is located within a highly urbanised landscape. Adjacent land to the south consists of cleared exotic parkland with patches of native vegetation. Isolated street trees exist within the surrounding residential areas. Given the lack of connectivity to large patches of habitat, only highly mobile threatened species are likely to occur within the development site (eg. Bats and birds).	Aerial imagery
Geological features of significance and soil hazard features	There are no geological features such as karst, caves, crevices or cliffs within the development site.	Aerial imagery
Areas of Outstanding Biodiversity Value	No areas of Outstanding Biodiversity Values occur within the development site.	Register of Declared Areas of Outstanding Biodiversity Value (DPIE 2020)
NSW (Mitchell) Landscapes	Cumberland Plain	NSW (Mitchell) Landscapes - version 3.1 (DPIE 2016)
Additional features required to be assessed	N/A. Any additional applicable legislation has been addressed in Section 8, including the EPBC Act.	
Percent (%) native vegetation extent	There are no differences between the mapped vegetation extent and the aerial imagery. The development footprint is approximately 0.27 ha and contains approximately 0.03 ha of native vegetation. The assessment area is approximately 755 ha and contains approximately 99 ha of native vegetation (13 %).	Calculated using aerial imagery and ArcGIS software

3. Native Vegetation

3.1. Survey Effort

Vegetation survey was undertaken within the development site by ELA ecologists Karen Spicer and Griffin Taylor-Dalton on 5 October 2021 (Figure 5).

One full-floristic vegetation plot was surveyed to identify Plant Community Types (PCTs) and Threatened Ecological Communities (TECs) on the development site (Table 3). The plot was undertaken on the development site to assess the composition, structure and function components of each vegetation zone in accordance with the BAM.

All field data collected at full-floristic and vegetation integrity plots is included in Appendix B and Appendix C.

Table 3: Full-floristic PCT identification plots

PCT ID	PCT Name	Number of plots surveyed
849	Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion	1

3.2. Native vegetation extent on the development site

The development site consists of cleared exotic park land vegetation, which is regularly mown, with patches of native vegetation present as canopy trees. There is no difference between the vegetation extent shown on the aerial imagery in this report and the extent of vegetation within the development site.

Under the BAM, the proposal has been assessed as a site-based development (i.e. not a linear development). Based on the amount of clearing proposed, this proposal has been assessed via Appendix C of the BAM: *Streamlined assessment module – small area*, as less than 1 ha of vegetation is proposed for removal.




Figure 3: Survey effort

3.3. Plant Community Types present

Plant community types (PCTs) identified within the development site are presented in Table 4 and Figure 4 . The development site also contained cleared areas, exotic vegetation and planted native vegetation which did not correspond with any PCT.

Table 4: Plant community types

PCT ID	PCT Name	Vegetation Class	Vegetation Formation	Area	Percent cleared
849	Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion	Coastal Valley Grassy Woodland	Grassy Woodlands	0.03ha	93%

3.3.1. Plant Community Type selection justification

In determining the PCT for the development site, various attributes were considered in combination to assign vegetation to the best fit PCT. Attributes included dominant species in each stratum and relative abundance, community composition, soils and landscape position. Reference was made to the PCT descriptions in the BioNet Vegetation Classification and the final scientific determinations for TECs. Possible PCT options are provided in Table 5.

Table 5: Potential PCTs

Selected PCT ID	PCT Name	Other PCT options
849	Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion	724, 725, 835, 850

- PCT 724 *Castlereagh shale - gravel transition forest* is associated with shale-influenced sandy soils that support a component of ironstone gravels. Typically, the canopy includes *Eucalyptus fibrosa* (Broad-Leaved Ironbark) along with a wide variety of other eucalypts depending on location and a lower canopy of *Melaleuca decora*. *Eucalyptus fibrosa* was recorded in low abundance within the development footprint. However, the presence of canopy species *E. tereticornis* and *E. molucanna* and clay soils align more with PCT 849.
- PCT 725 *Castlereagh Ironbark forest* is associated with clay soils derived from Tertiary alluvial deposits and has a typical structure of low dense thickets of paperbarks with low emergent eucalypts. The dominant canopy species is *Eucalyptus fibrosa* (Broad-Leaved Ironbark). However, the dominant species throughout the development site is *E. molucanna*, which is more aligned with PCT 849.
- PCT 835 *Cumberland riverflat forest* occurs on alluvial flats of the Cumberland Plain. The development site is located within 40 m of a first order drainage line, which may have supported vegetation consistent with PCT 835 prior to historical vegetation removal. As the development

site is elevated about this drainage line and is not located within a riverflat or low-lying drainage area, this PCT was disregarded.

- The landscape position within the relatively flat land of the Cumberland Plain eliminated PCT 850 *Cumberland Shale Hills Woodland*, which has similar canopy species as PCT 849, but occurs on higher elevations associated with the hills and rises south from Prospect (BioNet VIS, DPIE 2021).

3.4. Threatened Ecological Communities

PCT 849 is associated with the TEC *Cumberland Plain Woodland in the Sydney Basin Bioregion* listed as a Critically Endangered Ecological Community (CEEC) under the BC Act. PCT 849 is also associated with the TEC *Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest* listed as a CEEC under the EPBC Act. To be recognised under the EPBC Act, the vegetation must meet the condition thresholds outlined in the approved listing advice (DEWHA 2009). The patch of vegetation within the development site did not meet the condition thresholds under the EPBC Act. Justification for this decision is outlined in section 8.1 of this report. Figure 5 shows the extent of TEC within the development site.

Table 6: Threatened Ecological Communities within the development footprint

PCT ID	BC Act			EPBC Act		
	Listing status	Name	Area (ha)	Listing status	Name	Area (ha)
849	Critically Endangered	<i>Cumberland Plain Woodland in the Sydney Basin Bioregion</i>	0.03	Critically Endangered	<i>Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest</i>	0

3.5. Vegetation integrity assessment

3.5.1. Vegetation zones

One vegetation zone was identified on the development site (Table 7, Figure 6), being low condition of PCT 849 *Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion*. Descriptions of the vegetation zone is provided in Table 8.

Table 7: Vegetation zones and vegetation integrity survey plots collected on the development site

Vegetation Zone	PCT ID	PCT Name	Condition	Area (ha)	Patch Size	Vegetation Integrity Survey Plots required	Vegetation Integrity Survey Plots collected
1	849	Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion	Low	0.03	+101	1	1

Table 8: Description of the vegetation zones

849 - Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion				
Vegetation formation/class	Grassy Woodlands			
Conservation status	NSW BC Act Critically Endangered: Cumberland Plain Woodland in the Sydney Basin Bioregion EPBC Act Critically Endangered: Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest (although this patch does not meet the EPBC condition threshold)			
Description	This community occurs along the shale plains of the Sydney Basin region on elevations of less than 150m. Rainfall is restricted to a narrow band between 750 and 950 millimetres per annum. Canopy height is typically between 15-20m. Mid-storey species generally occupy a height of 2-5m.			
Characteristic canopy trees		<i>Eucalyptus moluccana</i>		
Characteristic mid-storey		No characteristic native mid-storey species were observed during the field survey, due to the previous disturbance that has occurred within the development site.		
Characteristic groundcovers		<i>Dichondra repens, Oxalis perennans</i>		
Mean native richness		6		
Exotic species / HTW cover		<i>Briza subaristata, Eragrostis curvula, Paspalum dilatatum, Senecio madagascariensis</i> – 71.2%		
Condition		Low condition		
Variation and disturbance		High level of disturbance due to high number of exotic species and regular mowing/slashing of the understorey.		
No. sites sampled		1		
Threatened flora species		No threatened flora species were identified during the field survey		
Fauna habitats		Could provide potential foraging habitat for typical peri-urban species of birds and mammals including <i>Trichosurus vulpecula</i> (Brush-tailed Possum) and highly mobile threatened bats.		
Composition	Structure	Function	Vegetation Integrity Score	
3.8	13.1	16.5	9.4	

849 - Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion



Approximately 0.03 ha of native vegetation was identified within the development footprint. Of this native vegetation, 0.003 ha has been planted for landscape purposes, and 0.03 ha has been mapped by ELA as remnant trees associated with *PCT 849 Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion*.

The 0.003 ha of planted native vegetation has been assessed against *Appendix D: Streamlined assessment module - planted native vegetation* of BAM 2020. The decision-making process for this module is presented below. The remaining vegetation within the development site was either exotic or cleared land dominated by exotic grasses.

3.5.2. Decision-making key for planted native vegetation (Appendix D BAM 2020)

NB: text in italics is copied directly from Appendix D (BAM 2020)

1: Does the planted native vegetation occur within an area that contains a mosaic of planted and remnant native vegetation and which can be reasonably assigned to a PCT known to occur in the same IBRA subregion as the proposal?

i. Yes..... The planted native vegetation must be allocated to the best fit PCT and the BAM must be applied.

ii. No..... Go to 2.

Justification:

Although the planted native vegetation occurred within proximity to the remnant native vegetation, the species *Eucalyptus microcorys* does not naturally occur on the Cumberland Plain.

2: Is the planted native vegetation:

a. *planted for the purpose of environmental rehabilitation or restoration under an existing conservation obligation listed in BAM Section 11.9(2.), and*

b. *the primary objective was to replace or regenerate a plant community type or a threatened plant species population or its habitat?*

i. ~~Yes The planted native vegetation must be assessed in accordance with Chapters 4 and 5 of the BAM.~~

ii. *No..... Go to 3.*

Justification:

The vegetation within the development site was planted for landscaping purposes, visual amenity and shade. The natural distribution of *Eucalyptus microcorys* does not include the Cumberland Plain landscape, within which the development site is situated.

3. Is the planted/translocated native vegetation individuals of a threatened species or other native species planted/translocated for the purpose of providing threatened species habitat under one of the following:

a. *a species recovery project*

b. *Saving our Species project*

c. *other types of government funded restoration project*

d. *condition of consent for a development approval that required those species to be planted or translocated for the purpose of providing threatened species habitat*

e. *legal obligation as part of a condition or ruling of court. This includes regulatory directed or ordered remedial plantings (e.g. Remediation Order for clearing without consent issued under the BC Act or the Native Vegetation Act)*

f. *ecological rehabilitation to re-establish a PCT or TEC that was, or is carried out under a mine operations plan, or*

g. *approved vegetation management plan (e.g. as required as part of a Controlled Activity Approval for works on waterfront land under the NSW Water Management Act 2000)?*

i. ~~Yes The planted native vegetation must be assessed in accordance with Chapters 4 and 5 of the BAM.~~

ii. *No..... Go to 4.*

Justification:

Planted native vegetation within the development site does not include any individuals of threatened species nor is the planting associated with any of the above conservation projects.

4. Was the planted native vegetation (including individuals of a threatened flora species) undertaken voluntarily for revegetation, environmental rehabilitation or restoration without a legal obligation to secure or provide for management of the native vegetation?

i. Yes..... Go to D.2 Assessment of planted native vegetation for threatened species habitat (the use of Chapters 4 and 5 of the BAM are not required to be applied).

ii. No..... Go to 5.

Justification:

No. The primary purpose of the plantings was for landscaping purposes. The planted native species are not consistent with a PCT from the locality.

5. Is the native vegetation (including individuals of a threatened flora species) planted for functional, aesthetic, horticultural or plantation forestry purposes? This includes examples such as: windbreaks in agricultural landscapes, roadside plantings (including street trees, median strips, roadside batters), landscaping in parks, gardens and sport fields/complexes, macadamia plantations or teatree farms?

i. Yes Go to D.2 Assessment of planted native vegetation for threatened species habitat (the use of Chapters 4 and 5 of the BAM are not required to be applied).

ii. No..... Go to 6.

Justification:

The planted native vegetation within the development site was planted for the purposes of landscaping and amenity within Chameleon Reserve.

3.5.3. Patch size

Patch size was calculated using available vegetation mapping for all patches of intact native vegetation on and adjoining the development site. Patch size was assigned to one of four classes (<5 ha, 5-24 ha, 25-100 ha or ≥100 ha). A patch size 97 ha was determined for the development site.

3.5.4. Assessing vegetation integrity

A vegetation integrity assessment using the BAM Calculator (BAMC) was undertaken and the results are outlined in Table 9.

Table 9: Vegetation integrity scores

Veg Zone	PCT ID	Condition	Area (ha)	Composition Condition Score	Structure Condition Score	Function Condition Score	Presence of Hollow bearing trees	Current vegetation integrity score
1	849	Low	0.03	3.8	13.1	16.5	No	9.4



Figure 4: Plant Community Types



Figure 5: Threatened Ecological Communities




Figure 6: Vegetation Zones and Plots

4. Threatened species

4.1. Ecosystem credit species

Ecosystem credit species predicted to occur within the development site are generated by the BAMC following the input of VI data and the PCTs identified within Chapter 3. Ecosystem credit species predicted to occur at the development site, their associated habitat constraints, geographic limitations and sensitivity to gain class is included in Table 10.

Table 10: Predicted ecosystem credit species

Species	Common Name	Habitat Constraints	Geographic limitations	Sensitivity to gain class	BC Act listing status	EPBC Act Listing status
<i>Anthochaera phrygia</i>	Regent Honeyeater (Foraging)	-	-	High	CE	CE
<i>Artamus cyanopterus cyanopterus</i>	Dusky Woodswallow	-	-	Moderate	V	Not Listed
<i>Callocephalon fimbriatum</i>	Gang-gang Cockatoo (Foraging)	-	-	Moderate	V	Not Listed
<i>Chthonicola sagittata</i>	Speckled Warbler	-	-	High	V	Not Listed
<i>Circus assimilis</i>	Spotted Harrier	-	-	Moderate	V	Not Listed
<i>Climacteris picumnus victoriae</i>	Brown Treecreeper	-	-	High	V	Not Listed
<i>Daphoenositta chrysoptera</i>	Varied Sitella	-	-	Moderate	V	Not Listed
<i>Dasyurus maculatus</i>	Spotted-tailed Quoll	-	-	High	V	E
<i>Glossopsitta pusilla</i>	Little Lorikeet	-	-	High	V	Not Listed
<i>Grantiella picta</i>	Painted Honeyeater	Mistletoes present at a density of greater than five mistletoes per hectare	-	Moderate	V	V
<i>Haliaeetus leucogaster</i>	White-bellied Sea-Eagle (Foraging)	Within 1km of a rivers, lakes, large dams or creek, wetlands and coastlines	-	High	V	Not Listed

Species	Common Name	Habitat Constraints	Geographic limitations	Sensitivity to gain class	BC Act listing status	EPBC Act Listing status
<i>Hieraaetus morphnoides</i>	Little Eagle (Foraging)	-	-	Moderate	V	Not Listed
<i>Hirundapus caudacutus</i>	White-throated Needletail	-	-	High	Not Listed	V
<i>Lathamus discolor</i>	Swift Parrot (Foraging)	-	-	Moderate	E	CE
<i>Lophoictinia isura</i>	Square-tailed Kite (Foraging)	-	-	Moderate	V	Not Listed
<i>Melanodryas cucullata cucullata</i>	Hooded Robin (south-eastern form)	-	-	Moderate	V	Not Listed
<i>Melithreptus gularis gularis</i>	Black-chinned Honeyeater (eastern subspecies)	-	-	Moderate	V	Not Listed
<i>Micronomus norfolkensis</i>	Eastern Coastal Free-tailed Bat	-	-	High	V	Not Listed
<i>Miniopterus australis</i>	Little Bent-winged Bat (Foraging)	-	-	High	V	Not Listed
<i>Miniopterus orianae oceanensis</i>	Large Bent-winged Bat (Foraging)	-	-	High	V	Not Listed
<i>Neophema pulchella</i>	Turquoise Parrot	-	-	High	V	Not Listed
<i>Ninox connivens</i>	Barking Owl (Foraging)	-	-	High	V	Not Listed
<i>Ninox strenua</i>	Powerful Owl (Foraging)	-	-	High	V	Not Listed
<i>Petroica boodang</i>	Scarlet Robin	-	-	Moderate	V	Not Listed
<i>Petroica phoenicea</i>	Flame Robin	-	-	Moderate	V	Not Listed
<i>Phascolarctos cinereus</i>	Koala (Foraging)	-	-	High	V	V
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox (Foraging)	-	-	High	V	V
<i>Saccolaimus flaviventris</i>	Yellow-bellied Sheath-tail-bat	-	-	High	V	Not Listed
<i>Stagonopleura guttata</i>	Diamond Firetail	-	-	Moderate	V	Not Listed

Species	Common Name	Habitat Constraints	Geographic limitations	Sensitivity to gain class	BC Act listing status	EPBC Act Listing status
<i>Tyto novaehollandiae</i>	Masked Owl (Foraging)	-	-	High	V	Not Listed

Justification for the inclusion or exclusion of ecosystem credit species can be found below in Table 11.

Table 11: Justification for exclusion of predicted ecosystem credit species

Species	Common Name	BC Act listing status	EPBC Act Listing status	Justification for exclusion of species
<i>Grantiella picta</i>	Painted Honeyeater	V	V	It is unlikely that the species would use the development site due to the absence of mistletoes within the site.

4.2. Species credit species

4.2.1. Identification of species credit species

As this BDAR is being assessed via the streamlined (small area) assessment (BAM 2020, Appendix C), only those candidate species at risk of a serious and irreversible impact (SAIL) or those recorded on site require targeted survey. Species credit species generated by the BAMC, their associated habitat constraints, geographic limitations and sensitivity to gain class is included in Table 12. All the candidate species generated by the BAM C are entities at risk of a SAIL.

Table 13 outlines the justification for the inclusion or exclusion of these species.

Data utilised to inform the below decisions was derived from a variety of sources, including but not limited to the Threatened Biodiversity Data Collection/BioNet Atlas Database (BioNet 2021) and the Important Areas Map on the Biodiversity Offsets and Agreements Management System (BOAMS 2021).

Table 12: Candidate species credit species

Species	Common Name	Habitat Constraints	Geographic limitations	Sensitivity to gain class	BC Act listing status	EPBC Act Listing status
<i>Anthochaera phrygia</i>	Regent Honeyeater (Breeding)	As per mapped areas	-	High	CE	CE
<i>Caladenia tessellata</i>	Thick Lip Spider Orchid	-	-	Moderate	E	V
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	Within two kilometres of rocky areas containing caves, overhangs, escarpments, outcrops, or crevices, or within two kilometres of old mines or tunnels	-	Very High	V	V

Species	Common Name	Habitat Constraints	Geographic limitations	Sensitivity to gain class	BC Act listing status	EPBC Act Listing status
<i>Lathamus discolor</i>	Swift Parrot (Breeding)	As per mapped areas	-	Moderate	E	CE
<i>Miniopterus australis</i>	Little Bent-winged Bat (Breeding)	Cave, tunnel, mine, culvert or other structure known or suspected to be used for breeding including species records in BioNet with microhabitat code 'IC – in cave' observation type code 'E nest-roost' with numbers of individuals >500 or from the scientific literature	-	Very high	V	Not listed
<i>Miniopterus orianae oceanensis</i>	Large Bent-winged Bat (Breeding)	Cave, tunnel, mine, culvert or other structure known or suspected to be used for breeding including species records with microhabitat code "IC - in cave" observation type code "E nest-roost" with numbers of individuals >500	-	Very high	V	Not listed

Table 13: Justification for the exclusion/ inclusion of species credit species

Species	Common Name	BC Act listing status	EPBC Act Listing status	Justification for exclusion of species
<i>Anthochaera phrygia</i>	Regent Honeyeater (Breeding)	CE	CE	Excluded: The development site is not mapped as an important area for the Regent Honey Eater.
<i>Caladenia tessellata</i>	Thick Lip Spider Orchid	E	V	Excluded: Currently, this species is only known to occur in populations near Braidwood on the NSW Southern Tablelands and around the Wyong Area along the NSW Central Coast. It has been suggested that this species requires frequent fire regimes to stimulate flowering and seeding events (Duncan 2010). Given the location of the development site, it is highly likely that it has been a long time since the development site has been burnt. Given the above information, and the degraded nature of the development site, it is highly unlikely that this species would occur.

Species	Common Name	BC Act listing status	Act	EPBC Listing status	Act	Justification for exclusion of species
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	V		V		Excluded: There are no identified rocky areas or old mines or tunnels located within two kilometres of the development site.
<i>Lathamus discolor</i>	Swift Parrot (Breeding)	E		CE		Excluded: The development site is not mapped as an important area for the Swift Parrot.
<i>Miniopterus australis</i>	Little Bent-winged Bat (Breeding)	V		Not listed		Excluded: There were no caves or culverts to utilise across the development site. As a result, it is very unlikely that this species would utilise the site for breeding habitat.
<i>Miniopterus orianae oceanensis</i>	Large Bent-winged Bat (Breeding)	V		Not listed		Excluded: There were no caves or culverts to utilise across the development site. As a result, it is very unlikely that this species would utilise the site for breeding habitat.

4.2.2. Assessment of habitat constraints and vagrant species

Habitat for candidate species derived by the BAM C are not considered likely to occur within the development site based on the lack of habitat and long-term habitat degradation throughout the development site (justification provided in

Table 13).

ELA Botanist and orchid expert Lachlan Copeland was consulted regarding the likelihood of *Caladenia tessellata* and the requirement for targeted surveys. Lachlan noted the orchid would not occur within the development site given the following:

- In the last 15 years *C. tessellata* has only been known from two locations in NSW, despite several historical records for Greater Sydney
- The species is mainly known from sandy soils and it would likely never have occurred on the fertile soils of the Cumberland Plain
- The species is sensitive to disturbance
- It mainly flowers the year after fire, so unless the development site was burnt the previous summer/early Autumn, surveying for this species, even in a good season in spring, would be futile
- The four locations the species is known from in Victoria / NSW are characterised by shallow sandy soils with a mostly shrubby ground-layer. Given the development site is within fertile clay / loam soils with a grassy understory, there is no habitat for this species.

No other candidate species (in addition to those generated by the BAM C) are considered likely to occur within the development site and no further assessment is required.

4.2.3. Candidate species requiring further assessment

As this BDAR is being assessed via the streamlined (small area) assessment (BAM 2020, Appendix C), only those candidate species at risk of a serious and irreversible impact (SAII) require targeted survey.

All the candidate species generated by the BAM C (Table 12) are entities at risk of a SAIL. However, as there is no breeding habitat present within the development site for the fauna species (*Anthochaera Phrygia*, *Chalinolobus dwyeri*, *Lathamus discolor*, *Miniopterus australis*, *Miniopterus oriana oceanensis*) and no habitat for *C. tessellata*, no further assessment or targeted survey is required.

4.3. Identification of prescribed additional biodiversity impact entities

4.3.1. Karst, caves, crevices, cliffs, rocks and other geological features of significance

There are no karst, caves, crevices, cliffs, rocks or other geological features of significance within the development site.

4.3.2. Human-made structures and non-native vegetation

There are no human-made structures within the development site that will require removal.

Non-native vegetation within the development footprint consists of ground cover species that do not provide habitat for threatened species.

4.3.3. Habitat connectivity

The development site is located within a highly urbanised landscape. Adjacent land to the north and west consists of cleared exotic parkland with patches of native vegetation and sporting fields. Other small reserves and isolated street trees exist within the surrounding residential areas. Given the lack of connectivity to large patches of habitat, only highly mobile threatened species are likely to occur within the development site (e.g. bats).

4.3.4. Water bodies, water quality and hydrological processes

There are no water bodies present within the development site.

4.3.5. Wind farm developments

Not applicable to this development.

4.3.6. Vehicle strikes

Given the locality is already highly urbanised, vehicle strike is an existing threat to fauna. The proposed development will not involve the construction of internal roads, and therefore no additional threat to fauna from vehicle strike, beyond the existing levels, is likely to result.

5. Avoiding and Minimising Impacts on Biodiversity Values

5.1. Locating a project to avoid and minimise impacts on biodiversity values

5.1.1. Direct and indirect impacts

The BC Act and BAM 2020 requires proponents to avoid and minimise impacts on biodiversity values (refer to Section 7.1.1 of the BAM 2020). The development site contains 0.08 ha of PCT849, mostly consisting of large *E. molucanna* trees to the west of the carparking area, however none of these trees will be impacted by the proposed development.

The construction of the residences, driveways and landscaped areas within the development footprint will require the removal of (0.03 ha) (Figure 4) of native vegetation, which includes 0.03 ha of PCT 849_poor and 0.003 ha of planted native vegetation. As such, 0.05 ha of PCT 849_poor will be retained within the development site. In addition, canopy overhanging the site from adjacent trees will not be impacted (Laws 2021).

As discussed in Section 3.4, this vegetation zone conforms to the TEC *Cumberland Plain Woodland in the Sydney Basin Bioregion*. This vegetation community is listed as Critically Endangered under the BC Act. Avoiding and minimising impacts to the patches of Cumberland Plain Woodland within the development site is unlikely to result in the long-term biodiversity gains for this TEC, given:

- the existing poor condition of the TEC within the development site, which consists of canopy species and a ground cover dominated by exotic species
- the lack of connectivity with large patches of intact native vegetation
- the highly urbanised nature of the locality is unlikely to provide for future expansion in the extent and/or restoration of the TEC, or for connectivity with other native vegetation to be established.

Apart from the removal of 0.03 ha of Cumberland Plain Woodland TEC, the proposed development is avoiding and minimising impacts on biodiversity as follows:

- the development footprint is positioned between an existing carpark and residential area, within an area of high exotic groundcover and a sparse upper-storey, minimising the extent of native vegetation removed
- a large proportion of the development footprint consists of cleared land or exotic vegetation with no to poor biodiversity values and does not require offsetting or assessment under the BAM
- there are no impacts to breeding habitat for threatened species and no requirement to offset impacts to Species Credit Species
- given the lack of habitat connectivity within the development site, available foraging habitat for ecosystem species included in the assessment (see Table 10) is marginal
- indirect impacts to adjacent habitat are unlikely to result from the proposed development given the adjacent land is highly disturbed, primarily consisting of gardens and cleared lawn
- impacts to adjacent trees within the development site has been assessed by Laws 2021, with no impacts to adjacent trees expected to result

- measures to minimise impacts to biodiversity will include supervision of vegetation removal by an ecologist to ensure any fauna present are taken into care and relocated into similar nearby habitat

5.1.2. Prescribed biodiversity impacts

Prescribed biodiversity impacts are discussed in Section 4.3. There are no relevant prescribed impacts to be avoided or minimised.

5.2. Designing a project to avoid and minimise impacts on biodiversity values

5.2.1. Direct and indirect impacts

The project has been designed consistent with surrounding development in the locality. The majority of the development footprint does not require assessment, as it is cleared land. Most of the vegetation within the development site will be retained and maintained for recreational purposes. The project has been designed to create two new residential areas consistent with the surrounding land use.

5.2.2. Prescribed biodiversity impacts

Prescribed biodiversity impacts are likely to be minimal and the measures above will ensure they are minimised.

6. Assessment of Impacts

6.1. Direct impacts

The direct impacts of the development on:

- native vegetation and threatened ecological communities are outlined in Table 14
- prescribed biodiversity impacts are outlined in Section 6.4.

Direct impacts including the final project footprint (construction and operation) are shown on Figure 7.

Table 14: Direct impacts to native vegetation

PCT ID	PCT Name	BC Act listing	EPBC Act listing	Direct impact (ha)
849	Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion	CE	N/A to the development site, as the condition of the vegetation does not meet EPBC Act thresholds	0.03

6.2. Change in vegetation integrity

The change in vegetation integrity as a result of the development is outlined in Table 15.

Table 15: Change in vegetation integrity

Veg Zone	PCT ID	Condition	Area (ha)	Current vegetation integrity score	Future vegetation integrity score	Change in vegetation integrity
1	849	Low	0.03	9.4	0	-9.4




Figure 7: Combined impact map

6.3. Indirect impacts

The indirect impacts of the development are outlined in Table 16. Indirect impact zones may include residential housing to the east, Chameleon Drive to the south and Chameleon Reserve to the north and west. Most habitat within this indirect impact zone is highly disturbed and includes exotic gardens and cleared land.

Table 16: Indirect impacts

Indirect impact	Description (nature, extent and frequency)	Biodiversity affected	Duration/ Timing	Consequence
inadvertent impacts on adjacent habitat or vegetation	Inadvertent impacts to adjacent habitat/vegetation are likely to be small-scale and infrequent given the development footprint boundary is bordered by an existing road, carpark and fencing. There may be minor impacts to the patch of native vegetation cover to the north of the development footprint.	Impacts to threatened species or their habitat are unlikely given the adjacent habitat is highly disturbed and dominated by exotic vegetation.	During earthworks and construction	Minor impact
Reduced viability of adjacent habitat due to edge effects	Edge effects are likely to be small-scale and infrequent given the disturbed nature of the adjacent habitat.	As above	During earthworks and construction	Minor impact
Reduced viability of adjacent habitat due to noise, dust or light spill	Adjacent habitat is already highly disturbed residential land. Indirect impacts from dust, noise and light spills will not reduce the viability of adjacent habitat.	As above	Long-term	No impact
transport of weeds and pathogens from the site to adjacent vegetation	Adjacent land is already highly disturbed and dominated by exotic species and landscaped gardens. There is a risk of new weed species being introduced during site development from new residents.	As above	Long-term	Minor impact
Increased risk of starvation or exposure and loss of shade or shelter	N/A	N/A	N/A	N/A

Indirect impact	Description (nature, extent and frequency)	Biodiversity affected	Duration/ Timing	Consequence
loss of breeding habitat	No breeding habitat for threatened species is likely to occur within the indirect impact area, given the adjacent land is highly disturbed	No impacts expected	Long-term	None expected
trampling of threatened species	No threatened flora species are likely to occur within the indirect impact area, given the adjacent land is highly disturbed	No impacts expected	Long-term	None expected
Inhibition of nitrogen fixation and increased soil salinity	N/A	N/A	N/A	N/A
Fertiliser drift	N/A	N/A	N/A	N/A
rubbish dumping	N/A	N/A	N/A	N/A
wood collection	N/A	N/A	N/A	N/A
removal and disturbance of rocks including bush rock	N/A	N/A	N/A	N/A
increase in predators	N/A	N/A	N/A	N/A
increase in pest animal populations	N/A	N/A	N/A	N/A
changed fire regimes	N/A	N/A	N/A	N/A
disturbance to specialist breeding and foraging habitat, e.g. beach nesting for shorebirds.	N/A	N/A	N/A	N/A
sedimentation and contaminated and/or nutrient rich run-off	There is the risk of sediment movement during earthworks and site development impacting adjacent land.	No threatened species are likely to be affected	Short-term during earthworks and construction	Minor impact

6.4. Prescribed biodiversity impacts

The development site has the prescribed biodiversity impacts as outlined in Table 17.

Table 17: Direct impacts on prescribed biodiversity impacts

Prescribed biodiversity impact	Description (Nature, extent and frequency)	Consequences	Justification	Additional information
Karst, caves, crevices, cliffs,	N/A – no geological features are present	N/A	N/A	N/A

Prescribed biodiversity impact	Description (Nature, extent and frequency)	Consequences	Justification	Additional information
rocks and other geological features of significance				
Human made structures or non-native vegetation	N/A - there are no human-made structures within the development site that will require removal. Non-native vegetation within the development footprint consists of ground cover species that do not provide habitat for threatened species.	N/A	N/A	N/A
Habitat connectivity	There is a lack of connectivity to large patches of habitat, so only highly mobile threatened species are likely to occur within the development site (e.g. bats).	Minor loss of foraging habitat for threatened bat species that may forage across site including <i>Micronomus norfolkensis</i> (Eastern Coastal Free-tailed Bat), <i>Miniopterus australis</i> (Little Bent-winged Bat), <i>Miniopterus orianae oceanensis</i> (Large Bent-winged Bat) and <i>Pteropus poliocephalus</i> (Grey-headed Flying-fox).	Habitat for these threatened bats within the development site is marginal and is not well connected with areas of large intact foraging habitat for these species. Threatened bird species listed as predicted ecosystem species in Table 11 are unlikely to regularly occur within the development site, given the poor habitat condition (lack of structural and floral diversity) and lack of connectivity with large areas of native bushland.	N/A
Water bodies, water quality and hydrological processes	N/A - there are no water bodies present within the development site.	N/A	N/A	N/A
Wind turbine strikes on protected animals	N/A	N/A	N/A	N/A
Vehicle strikes	The locality is already highly urbanised, and vehicle strike is an existing threat to fauna. The proposed	N/A	N/A	N/A

Prescribed biodiversity impact	Description (Nature, extent and frequency)	Consequences	Justification	Additional information
	development does not involve construction of internal roads. Additional threats to fauna from vehicle strike beyond the existing levels, is unlikely to result.			

6.5. Mitigating and managing direct and indirect impacts

Measures proposed to mitigate and manage impacts at the development site before, during and after construction are outlined in Table 18.

Table 18: Measures proposed to mitigate and manage impacts

Measure	Risk before mitigation	Risk after mitigation	Action	Outcome	Timing	Responsibility
timing works to avoid critical life cycle events such as breeding or nursing	N/A	N/A	N/A	N/A	N/A	N/A
instigating clearing protocols including pre-clearing surveys, daily surveys and staged clearing, the presence of a trained ecological or licensed wildlife handler during clearing events	Low – no hollow-bearing trees, nests or dreys were recorded within the development site.	Low	An ecologist be present to supervise vegetation removal and care and relocation of fauna (if present)	Ensuring animal welfare and protection	During vegetation removal	Project ecologist
installing artificial habitats for fauna in adjacent retained vegetation and habitat or human made structures to replace the habitat resources lost and encourage animals to move from the impacted site, e.g. nest boxes	N/A – no important habitat identified within the development area.	Low	N/A	N/A	N/A	N/A

Measure	Risk before mitigation	Risk after mitigation	Action	Outcome	Timing	Responsibility
clearing protocols that identify vegetation to be retained, prevent inadvertent damage and reduce soil disturbance; for example, removal of native vegetation by chain-saw, rather than heavy machinery, is preferable in situations where partial clearing is proposed	N/A – the entire site will be cleared of vegetation	N/A	N/A	N/A	N/A	N/A
sediment barriers or sedimentation ponds to control the quality of water released from the site into the receiving environment	Moderate	Low	Standard sediment and erosion control (SEC) measures should be implemented	Contain sediments within the site	Establish measures prior to commencing works	SEC Site manager
noise barriers or daily/seasonal timing of construction and operational activities to reduce impacts of noise	N/A	N/A	N/A	N/A	N/A	N/A
light shields or daily/seasonal timing of construction and operational activities to reduce impacts of light spill	N/A	N/A	N/A	N/A	N/A	N/A
adaptive dust monitoring programs to control air quality	Moderate	Low	Standard sediment and erosion control (SEC) measures and dust control measures should be implemented	Control dust during dry periods	Watering of stockpiles exposed during periods	of and soil dry Site manager

Measure	Risk before mitigation	Risk after mitigation	Action	Outcome	Timing	Responsibility
programming construction activities to avoid impacts; for example, timing construction activities for when migratory species are absent from the site, or when particular species known to or likely to use the habitat on the site are not breeding or nesting	N/A	N/A	N/A	N/A	N/A	N/A
temporary fencing to protect significant environmental features such as riparian zones	N/A	N/A	N/A	N/A	N/A	N/A
hygiene protocols to prevent the spread of weeds or pathogens between infected areas and uninfected areas	Moderate	Low	Come clean – go clean protocol for plant and equipment entering/exiting the site	Reduce the risk of introducing new weed species	During earthworks and construction of subdivision	Site manager
staff training and site briefing to communicate environmental features to be protected and measures to be implemented	Moderate	Low	Standard SEC measures and come clean – go clean protocol.	Reduce the risk of introducing new weed species	Prior to commencement of earthworks	Site manager
development control measures to regulate activity in vegetation and habitat adjacent to residential development including controls on pet ownership, rubbish disposal, wood collection, fire management and disturbance to	N/A	N/A	N/A	N/A	N/A	N/A

Measure	Risk before mitigation	Risk after mitigation	Action	Outcome	Timing	Responsibility
nests and other niche habitats						
making provision for the ecological restoration, rehabilitation and/or ongoing maintenance of retained native vegetation habitat on or adjacent to the development site	N/A	N/A	N/A	N/A	N/A	N/A

6.6. Mitigating prescribed impacts

Measures proposed to mitigate and manage prescribed biodiversity impacts at the development site before, during and after construction are outlined in Table 19.

Table 19: Mitigation measures for prescribed biodiversity impacts

Measure	Risk before mitigation	Risk after mitigation	Action	Outcome	Timing	Responsibility
Scheduling timing of construction activities to avoid critical life cycle events	N/A	N/A	N/A	N/A	N/A	N/A
Instigating clearing protocols including pre-clearing surveys, daily surveys and staged clearing, and using a trained ecological or licensed wildlife handler during clearing, construction and maintenance activities for human made structures and non-native vegetation	N/A	N/A	N/A	N/A	N/A	N/A
Retaining habitat features within the development footprint or relocating them to adjacent retained remnant vegetation	N/A	N/A	N/A	N/A	N/A	N/A
Installing artificial connectivity measures to re-establish	N/A	N/A	N/A	N/A	N/A	N/A

Measure	Risk before mitigation	Risk after mitigation	Action	Outcome	Timing	Responsibility
connections between habitat and favoured transport corridors						
Erecting temporary fencing to protect significant environmental features such as karst, caves, rock outcrops and water bodies	N/A	N/A	N/A	N/A	N/A	N/A
Replacing habitat provided by human made structures and non-native vegetation with alternative habitat	N/A	N/A	N/A	N/A	N/A	N/A
Sediment barriers or sedimentation ponds to control the quality of water released from the site into the receiving environment	Moderate	Low	Standard sediment and erosion control (SEC) measures and dust control measures should be implemented	Contain sediments within the site to avoid entering stormwater	Establish SEC measures prior to works commencing	Site foreman
Staff training and site briefing to communicate environmental features to be protected and measures implemented to protect them	N/A	N/A	N/A	N/A	N/A	N/A
Ecological restoration, rehabilitation actions and/or maintenance of retained native vegetation on or adjacent to the development footprint	N/A	N/A	N/A	N/A	N/A	N/A
Development control measures that regulate the types of activities that can occur in native vegetation and habitat adjacent to residential development including prohibiting the collection of bush rocks	N/A	N/A	N/A	N/A	N/A	N/A

6.7. Adaptive management strategy

Adaptive management for uncertain biodiversity impacts (that are infrequent or difficult to measure) is not relevant or required for the proposed development.

7. Impact summary

Following implementation of the BAM and the BAMC, the following impacts have been determined.

7.1. Serious and Irreversible Impacts (SAII)

The development has candidate Serious and Irreversible Impacts (SAII) values as outlined in Table 20. Detailed consideration of whether impacts on TECs that are serious and irreversible is included in Table 21.

Table 20: Serious and Irreversible Impacts Summary

Ecological Community	Principle	Direct impact individuals / area (ha)	Threshold
Cumberland Woodland in the Sydney Bioregion	Plain in the Basin Principle 1 (species or ecological community currently in a rapid rate of decline), and Principle 2 (species or ecological communities with a very small population size)	0.03	Under development

Table 21: Evaluation of an impact on a TEC consistent with 9.1.1 of the BAM

Impact Assessment Provisions	Assessment
1. the action and measures taken to avoid the direct and indirect impact on the potential entity for an SAI	Measures taken to avoid the direct and indirect impacts to Cumberland Plain Woodland as addressed in Section 5. However, given the relatively small size of the development site and the highly disturbed native of the vegetation within the development site, all vegetation will be removed.
2a. evidence of reduction in geographic distribution (Principle 1, clause 6.7(2)(a) BC Regulation) as the current total geographic extent of the TEC in NSW AND the estimated reduction in geographic extent of the TEC since 1970 (not including impacts of the proposal)	Prior to European settlement, Cumberland Plain Woodland was extensive across the Cumberland Plain in western Sydney on soils derived from Wianamatta Shale (Bionet 2021), covering approximately 125,000 ha (NPWS 2004). The current geographic extent of Cumberland Plain Woodland is estimated to be 9% of this original extent (Bionet, 2021). The estimated reduction in geographic extent since 1970 was not found in relevant literature.
2b. extent of reduction in ecological function for the TEC using evidence that describes the degree of environmental degradation or disruption to biotic processes (Principle 2, clause 6.7(2)(b) BC Regulation) indicated by: i. change in community structure ii. change in species composition iii. disruption of ecological processes iv. invasion and establishment of exotic species v. degradation of habitat, and vi. fragmentation of habitat	Cumberland Plain Woodland was originally cleared for agriculture during European settlement given its fertile soils, gentle slopes and proximity to early settlement within Sydney (DECCW 2010). As a result, by the middle of the 19th century, most of the Cumberland Plain was being grazed or under cultivation. In more recent years, Cumberland Plain Woodland has been subject to clearing for industrial and residential development and for infrastructure including roads, to service Sydney’s growing population (DECCW 2010). In relation to the ecological function of the community: ii. The structure of the community has changed, primarily to a grassland where canopy and shrub species have been removed. In some areas, such grassland is still dominated by native grasses and is considered to be Derived Native Grassland (DNG),

Impact Assessment Provisions

Assessment

- particularly in areas where pasture improvement and application of fertiliser has not occurred. In other areas, primarily in rural land-use, under-scrubbing has occurred, where a canopy is retained, but the shrub and groundcover removed to allow for grazing and hobby farms. As such, the structure of the TEC has been simplified.
- iii. The **composition** of the community has changed with a general reduction in diversity of native species. As woodland has become fragmented and patch size has reduced, weed invasion has reduced diversity, particularly with long established weeds such as *Olea europea* subsp. *cuspidata* (African Olive). This species has thrived in remnant and regenerating patches of Cumberland Plain Woodland and displaces native species, reducing native flora diversity. Likewise, the diversity of fauna that originally inhabited Cumberland Plain Woodland has been reduced and simplified, with local extinctions of mammals and birds, particularly since the 1950s (DECCW 2010). In turn, some opportunistic native species have increased in numbers over time, including the Noisy Miner (*Manorina melanocephala*).
 - iv. **Ecological processes** are the interactions between biota and abiotic components such as water cycle, mineral cycle, decomposition and nutrient cycles, reproduction / pollination and food webs. With a reduction in the extent of Cumberland Plain Woodland into smaller isolated and fragmented patches and a reduction in biodiversity of flora and fauna, such ecological processes have been disrupted. Understanding these processes can be complex, but as an example, the loss of certain pollination vectors is likely to lead of local extinction of some flora species. Changes to fire regimes, particularly due to the urban setting of western Sydney is another example of disruptions to ecological processes.
 - v. As discussed above, **invasion and establishment of exotic species** has become a significant problem within Cumberland Plain Woodland. This community is particularly vulnerable to weed invasion due its grassy understorey, comparatively fertile soils and past agricultural landuses (DECCW 2010).
 - vi. The **clearing and fragmentation** of Cumberland Plain Woodland into smaller and more isolated patches makes the remaining of areas of Cumberland Plain Woodland more susceptible to habitat degradation. Degradation results from all the factors listed above including edge effects, weed

Impact Assessment Provisions	Assessment
	<p>invasion, disruption of ecological processes and loss of biodiversity.</p> <p>vii. Fragmentation of Cumberland Plain Woodland has reduced the resilience of the community. Larger remnants are usually more diverse, less susceptible to edge effects and catastrophic events like bushfire and the expected impacts of climate change (DECCW 2010).</p>
<p>2c. evidence of restricted geographic distribution (Principle 3, clause 6.7 (2) (c) BC Regulation), based on the TECs geographic range in NSW according to the:</p> <p>i. extent of occurrence</p> <p>ii. area of occupancy, and</p> <p>iii. number of threat-defined locations.</p>	<p>This principle does not apply to Cumberland Plain Woodland TEC. However, the distribution of Cumberland Plain Woodland TEC is restricted to the Cumberland Plain within the Sydney Basin Bioregion and originally occupied 125,211 ha (pre-1750) (DECCW 2010).</p>
<p>2d. evidence that the TEC is unlikely to respond to management (Principle 4, clause 6.7 (2) (d) BC Regulation).</p>	<p>This principle does not apply to Cumberland Plain Woodland TEC. The recovery plan (DECCW 2010) discusses how Cumberland Plain Woodland responds to recovery actions, including weed control and revegetation.</p>
<p>3. Where the TBDC indicated that data is ‘unknown’ or ‘data deficient’ for a TEC for a criterion listed in subsection 9.1.1(2), the assessor must record this in the BDAR or BCAR.</p>	<p>N/A</p>
<p>4a. the impact on the geographic extent of the TEC (Principles 1 and 3) by estimating the total area of the TEC to be impacted by the proposal:</p> <p>i. in hectares, and</p> <p>ii. as a percentage of the current geographic extent of the TEC in NSW.</p>	<p>The total area of Cumberland Plain Woodland TEC affected by the proposed development is 0.03 ha.</p> <p>Assuming that 9% of the original 125,449 ha of Cumberland Plain Woodland remains today (11,290 ha), the Cumberland Plain Woodland affected by the proposed development represents 0.00027% of the current geographic extent of the TEC in NSW.</p>
<p>4b. the extent that the proposed impacts are likely to contribute to further environmental degradation or the disruption of biotic processes (Principle 2) of the TEC by:</p> <p>i. estimating the size of any remaining, but now isolated, areas of the TEC; including areas of the TEC within 500 m of the development footprint or equivalent area for other types of proposals</p> <p>ii. describing the impacts on connectivity and fragmentation of the remaining areas of TEC measured by:</p> <ul style="list-style-type: none"> • distance between isolated areas of the TEC, presented as the average distance if the remnant is retained AND the average distance if the remnant is removed as proposed, and • estimated maximum dispersal distance for native flora species characteristic of the TEC, and • other information relevant to describing the impact on connectivity and fragmentation, such as the area to perimeter ratio for remaining areas of the TEC as a result of the development 	<p>The extent that the proposed development is likely to contribute to further environmental degradation and disruption of biotic processes of Cumberland Plain Woodland is discussed below, in relation to:</p> <p>i. Remaining areas of TEC within 500 m of the development footprint is 4.34 ha as previously mapped by OEH (2013) and 0.43 ha as validated by ELA (Figure 8), providing a total area of 4.77 ha. This figure reflects the highly urbanised landscape throughout Erskine Park. Each patch within the 500 m buffer of the development site is small and isolated.</p> <p>ii. The existing condition of the Cumberland Plain Woodland TEC within the development site is already highly fragmented. The vegetation is in a poor condition and consists of canopy species, no shrub layer and a ground cover dominated by exotic species/lawn. The vegetation within the development site is part of a larger patch within Chameleon Reserve that extends to the north and west. The removal of 0.03 ha of Cumberland Plain Woodland within the development</p>

Impact Assessment Provisions	Assessment
<p>iii. describing the condition of the TEC according to the vegetation integrity score for the relevant vegetation zone(s) (Section 4.3). The assessor must also include the relevant composition, structure and function condition scores for each vegetation zone.</p>	<p>footprint will marginally increase the distance between remaining patches of woodland (by 5 m) as shown in Figure 8.</p> <p>iii. Dispersal of native flora species characteristic of the TEC may be several kilometres when considering pollen vectors such as <i>Pteropus poliocephalus</i> and seed dispersal by wind and fauna. However, given the location of the development site is within a largely suburban landscape, dispersal and exchange of flora species characteristic of the Cumberland Plain Woodland is likely to be very limited.</p> <p>iv. The vegetation integrity score for the vegetation zone PCT849_low, which is the only vegetation zone within the development site containing the Cumberland Plain Woodland TEC, is 9.4. This score is derived from a composition score of 3.8, a structure score of 13.1, and a function score of 16.5. A breakdown of these scores is contained in Appendix C. These scores are low when compared to the benchmarks for the PCT 849 (BioNet VIS, DPIE 2021). For example, the shrub richness and cover were one and 0.1, but benchmark is 8 and 16 respectively. Tree richness was 1, but benchmark is 5; grass richness was 2 but benchmark is 12. The tree cover, of 20, was below the benchmark of 53. In terms of function data, there was no large tree within the plot, compared to benchmark of 3. Fallen log total length was 0 m within the plot, but benchmark is 40 m. In summary, the Cumberland Plain Woodland within the development footprint is a highly simplified patch lacking structural and floral diversity with a high level of weed invasion (71.2 score for High Threat Exotic).</p>

7.2. Impacts requiring offsets

There are no impacts from the development requiring offsets for native vegetation (ecosystem credits) or species credit species or their habitat Figure 7. As discussed previously, the vegetation integrity score was too low to require offsets and there is no breeding habitat for species credit species within the development footprint.

7.3. Impacts not requiring offsets

The impacts of the development not requiring offset for native vegetation are outlined in Table 22 and shown on Figure 7.

Table 22: Impacts to native vegetation that do not require offsets

Vegetation Zone	PCT ID	PCT Name	Direct impact (ha)	Rationale
1	849_low	Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion	0.03	The vegetation integrity score calculated by the BAM C was 9.4, which is too low to require offsets

Impacts to native planted vegetation (0.003 ha) does not require offsetting, as demonstrated in Section 3.5.2.

7.4. Areas not requiring assessment

Areas not requiring assessment are shown in Figure 7. These areas do not contain native vegetation and do not need to be assessed for ecosystem credits. These areas have been assessed for threatened species habitat and prescribed impacts.

7.5. Credit summary

The number of ecosystem credits required for the development are zero as outlined in Table 23. There are no species credits required for this development. A biodiversity credit report is included in Appendix D.

Table 23: Ecosystem credits required

Vegetation Zone	PCT ID	PCT Name	Credit Class	Direct impact (ha)	Credits required
1	849_low	Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion	Cumberland Plain Woodland in the Sydney Basin Bioregion	0.03	0




Figure 8: Serious and Irreversible Impacts

8. Consistency with legislation and policy

Additional matters relating to impacts on flora and fauna which are not covered by the BC Act must also be addressed for the proposed development. Potential Matters of National Environmental Significance (MNES) in accordance with the EPBC Act have been addressed in Section 8.1. The likelihood of occurrence table for the EPBC Act listed species is provided in Appendix E.

8.1. *Environment Protection and Biodiversity Conservation Act 1999 (EPBC Act)*

The EPBC Act establishes a process for assessing the environmental impact of activities and developments where Matters of National Environmental Significance (MNES) may be affected. Under the Act, any action which 'has, will have, or is likely to have a significant impact on a matter of MNES' is defined as a controlled action, and requires approval from the Commonwealth Department of Agriculture Water and Environment (DAWE), which is responsible for administering the EPBC Act.

The process includes undertaking an Assessment of Significance for listed threatened species and ecological communities that represent a matter of MNES that will be affected as a result of the proposed action. Significant impact guidelines that outline a number of criteria have been developed by the Commonwealth of Australia (2013), to provide assistance in conducting the Assessment of Significance and help decide whether or not a referral to the Commonwealth is required.

8.1.1. Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest ecological community

Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest ecological community is listed as Critically Endangered under the EPBC Act. This TEC is confined to the Sydney Basin. Since European settlement, it has been estimated that the total extent of this vegetation has been reduced by 93% (BioNet Atlas, DPIE 2021).

Within the development site, no patches of PCT 849 vegetation met the condition thresholds outlined in the approved listing advice (Commonwealth of Australia, 2010). The flowchart shown in Figure 9 demonstrates why the vegetation does not meet the EPBC condition thresholds for this community. As PCT 849 vegetation is not recognised under the EPBC Act, a test of significance under the EPBC Act is not required.

Flowchart of key diagnostic features and condition thresholds to identify the Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest ecological community




Figure 9: Flowchart to assist with the determination of EPBC Act condition thresholds (Commonwealth of Australia, 2010)

8.1.2. *Pteropus poliocephalus* (Grey-headed Flying-fox)

The Grey-headed Flying-fox is listed as a vulnerable species under the EPBC Act. This species utilises a wide variety of habitats (including disturbed areas) for foraging and have been recorded travelling long distances on feeding forays. Fruits and flowering plants of a wide variety of species are the main food source. The species roosts in large 'camps' of up to 200,000 individuals. Camps are usually formed close to water and along gullies, however, the species has been known to form camps in urban areas.

The closest camps are listed below:

- Ropes Creek – approximately 3 km to the north
- Wetherill Park – approximately 11 km to the east.

There is potential that this species could use the development site on occasion for foraging purposes. The vegetation within the development footprint provides marginal foraging habitat. This includes the 0.03 ha of PCT 849 and 0.003 ha of planted native vegetation. An EPBC Act impact assessment has been undertaken for this species. The result of this assessment can be found in Table 24.

Table 24: EPBC Act impact assessment for *Pteropus poliocephalus* (Grey-headed Flying-fox)

Criterion	Assessment
Criterion a: lead to a long-term decrease in the size of an important population of a species	<p>The Matters of National Environmental Significance Impact Guidelines 1.1 (Commonwealth of Australia, 2013) defines an important population as a population that is necessary for a species' long-term survival and recovery. This may include populations identified as such in recovery plans, and/or that are:</p> <ul style="list-style-type: none"> • Key source populations either for breeding or dispersal • Populations that are necessary for maintaining genetic diversity, and/or • Populations that are near the limit of the species range <p>The development site does not provide habitat for a breeding camp however it could provide marginal foraging habitat for travelling Flying-foxes. Considering the small size of the impact and the amount of more suitable foraging habitat in the locality, it is considered unlikely that action will result on a long-term decrease in the size of an important population.</p>
Criterion b: reduce the area of occupancy of an important population	<p>Considering the small size of the impact and suitable areas of occupancy in the locality, it is considered unlikely that the proposed development will result in a long-term reduction of an area of occupancy of an important population.</p>
Criterion c: fragment an existing important population into two or more populations	<p>The potential seasonal foraging habitat to be removed is considered marginal relative to nearby potential foraging habitat within the locality. While the potential foraging habitat may contribute as a 'steppingstone' for this highly mobile species to other more substantial foraging habitat sites, this function is unlikely to be significantly inhibited by the proposed works, given that scattered vegetation and landscaped gardens will remain in the locality. This species has been recorded in urban environments and is likely to continue to forage adjacent to the development site within Chameleon Reserve and across the broader locality. Therefore, the proposed works are unlikely to fragment an existing important population into two or more populations.</p>
Criterion d: adversely affect habitat critical to the survival of a species	<p>Foraging habitat critical to the survival of this species includes natural habitat that is productive during winter and spring when food bottlenecks have been identified. Critical foraging habitat has also been known to support populations of > 30,000 individuals within a 50km radius, productive during the final stages of gestation, during the weeks of birth, lactation and conception (September to May), productive during the final stages of fruit development and ripening in commercial crops affected by Grey-headed Flying-fox, and known to support a continuously occupied camp (DECCW 2009). The vegetation to be removed is a small amount of</p>

Criterion	Assessment
	potential foraging resources (0.03 ha) in the wider region. Given that this species travels up to 50 km to forage, it is considered unlikely that the works would adversely affect habitat critical to the survival of this species.
Criterion e: disrupt the breeding cycle of an important population	The nearest active Grey-headed Flying-fox camp occurs approximately 3 km to the north of the development site at the Ropes Creek (DAWE 2021). Considering this distance, the marginal nature of the foraging resources in the development site, and the amount of other suitable foraging resources in the locality, it is considered unlikely that the action will disrupt the breeding cycle of an important population.
Criterion f: modify, destroy, remove or isolate or decrease the availability or quality of habitat to the extent that the species is likely to decline	Given the small amount of potential foraging habitat to be removed relative to the amount remaining in the locality, it is considered that potential foraging habitat will persist adjacent to the development site and across the locality. This species is highly mobile, and it is unlikely that the habitat to be removed would cause the species to decline. Furthermore, according to the National Flying-fox Monitoring Program, no Grey-headed Flying-fox camps currently occur or have ever been recorded within the development site (DAWE 2021). The nearest active Grey-headed Flying-fox camp occurs approximately 3 km to the north of the development site at the Ropes Creek camp (DAWE 2021). Therefore, no known Grey-headed Flying-fox roosting camps for this species will be affected by the proposed works.
Criterion g: Result in invasive species that are harmful to a vulnerable species becoming established in the vulnerable species' habitat	The proposed works will not result in the establishment of an invasive species that is harmful to Grey-headed Flying-fox.
Criterion h: Introduce disease that may cause the species to decline	The proposed works will not result in the introduction of a disease that is harmful to the Grey-headed Flying-fox.
Criterion i: Interfere substantially with the recovery of the species	Considering the above factors, the proposed works will not interfere substantially with the recovery of the species.
Conclusion	In consideration of the above, the proposed works are considered unlikely to have a significant impact on the Grey-headed Flying-fox. No camps are located closer than 3 km away from the development site. The removal of 0.03 ha of native vegetation is unlikely to affect the availability of foraging habitat for this species. The vegetation to be affected is similar in composition to the surrounding environment, as a result foraging Grey-headed Flying-fox will have ample choice of habitat should they forage in this area. As such, a referral is not recommended.

9. Conclusion

Eco Logical Australia Pty Ltd (ELA) was engaged by Penrith City Council to prepare a Streamlined (small area) Biodiversity Development Assessment Report (BDAR) for Lot 1106 DP709078 and Lot 1107 DP709078, Chameleon Dr, Erskine Park (the development site) within the Penrith local government area. The proposed development includes subdivision of Chameleon Reserve into five lots and development of four residential lots with a development footprint of 0.27 ha.

Native vegetation was mapped within the development site, consisting of 0.003 ha of planted native vegetation and 0.03 ha of PCT 849 *Grey Box - Forest Red Gum grassy woodland on flats of the Cumberland Plain, Sydney Basin Bioregion*. PCT 849 was present in one condition state – low. The remaining vegetation within the development site was either exotic or cleared land dominated by exotic grasses.

The 0.003 ha of planted native vegetation was assessed against Appendix D: *Streamlined assessment module - planted native vegetation* of the BAM 2020 and does not require offsetting. The removal of 0.03 ha of PCT 849_low does not require offsets. No species credits are required to offset impacts.

PCT 849 corresponds to the threatened ecological community *Cumberland Plain Woodland in the Sydney Basin Bioregion*. This TEC is also an entity at risk of Serious and Irreversible Impact (SAIL) and an assessment has been undertaken to assist the consent authority to decide whether a SAIL will result.

Supervision of vegetation removal by an ecologist should be undertaken to ensure any fauna present are taken into care and relocated into similar nearby habitat. Sediment and erosion control measures are recommended to prevent stormwater pollution or movement of soils from the development site.

10. References

Biodiversity Offsets and Agreements Management System (BOAMS 2021).

<https://customer.lmbc.nsw.gov.au/assessment/s/userlogin> (logged in to access the BAM calculator, 6 December 2021)

Commonwealth of Australia. 2013. *Matters of National Environmental Significance - Significant impact guidelines 1.1 Environment Protection and Biodiversity Conservation Act 1999*. Prepared by the Department of the Environment.

Department of Agriculture Water and Environment (DAWE) 2021. *National Flying-fox monitoring viewer*. Australian Government. URL: <http://www.environment.gov.au/webgis-framework/apps/ffc-wide/ffc-wide.jsf>.

Department of Agriculture Water and Environment (DAWE). 2021. Protected Matters Search Tool. URL: <http://www.environment.gov.au/epbc/protect/index.html>

Department of Environment, Climate Change and Water NSW (DECCW) 2009. *Draft National Recovery Plan for the Grey-headed Flying-fox Pteropus poliocephalus*. Prepared by Dr Peggy Eby. Department of Environment, Climate Change and Water NSW, Sydney

Department of Environment, Climate Change and Water NSW (DECCW) 2010. *Cumberland Plain Recovery Plan*. Department of Environment, Climate Change and Water (NSW), Sydney.

Department of Environment, Water, Heritage and the Arts NSW (DEWHA) 2009. *Approved Conservation Advice for Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest ecological community*.

DEWHA 2010. *Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest, A guide to identifying and protecting the nationally threatened ecological community Environment Protection and Biodiversity Conservation Act 1999, Policy Statement 3.31*. Australian Government, Department of the Environment, Water, Heritage and the Arts.

Department of Planning, Industry and Environment (DPIE) 2021. BioNet Threatened Biodiversity Data Collection. URL: <http://www.bionet.nsw.gov.au/>. Accessed 29 September 2021

Department of Planning, Industry and Environment (DPIE) 2021. BioNet VIS Vegetation Classification URL: <https://www.environment.nsw.gov.au/research/Visclassification.htm>.

Department of Planning, Industry and Environment (DPIE) 2021. Biodiversity Values Map and Threshold Tool <https://www.lmbc.nsw.gov.au/Maps/index.html?viewer=BOSETMap>

Department of Planning, Industry and Environment (DPIE) 2021. ePlanning Spatial Viewer <https://www.planningportal.nsw.gov.au>

Laws, G. 2021. Arboricultural Impact Assessment Report & Tree Management Plan. Prepared for Penrith City Council, November 2021.

National Parks and Wildlife Service (NPWS) (NSW) 2004. Cumberland Plain Woodland, Endangered Ecological Community Information.

OEH 2013. *Remnant Vegetation of the western Cumberland subregion*, 2013 Update. VIS_ID 4207

Appendix A Definitions

The following terminology has been used throughout this report for the purposes of describing the impacts of the proposal in the context of a biodiversity assessment in accordance with the NSW Biodiversity Assessment Method 2020. This terminology may or may not align with other technical documents associated with the proposed development.

Terminology	Definition
Biodiversity credit report	The report produced by the Credit Calculator that sets out the number and class of biodiversity credits required to offset the remaining adverse impacts on biodiversity values at a development site, or on land to be biodiversity certified, or that sets out the number and class of biodiversity credits that are created at a biodiversity stewardship site.
BioNet Atlas	The BioNet Atlas (formerly known as the NSW Wildlife Atlas) is the DPIE database of flora and fauna records. The Atlas contains records of plants, mammals, birds, reptiles, amphibians, some fungi, some invertebrates (such as insects and snails) and some fish
Broad condition state:	Areas of the same PCT that are in relatively homogenous condition. Broad condition is used for stratifying areas of the same PCT into a vegetation zone for the purpose of determining the vegetation integrity score.
Connectivity	The measure of the degree to which an area(s) of native vegetation is linked with other areas of vegetation.
Credit Calculator	The computer program that provides decision support to assessors and proponents by applying the BAM, and which calculates the number and class of biodiversity credits required to offset the impacts of a development or created at a biodiversity stewardship site.
Development	Has the same meaning as development at section 4 of the EP&A Act, or an activity in Part 5 of the EP&A Act. It also includes development as defined in section 115T of the EP&A Act.
Development footprint	The area of land that is directly impacted on by a proposed development, including access roads, and areas used to store construction materials.
Development site	An area of land that is subject to a proposed development that is under the EP&A Act.
Ecosystem credits	A measurement of the value of EECs, CEECs and threatened species habitat for species that can be reliably predicted to occur with a PCT. Ecosystem credits measure the loss in biodiversity values at a development site and the gain in biodiversity values at a biodiversity stewardship site.
Extent of occurrence (EOO)	Measures the spatial spread of a taxon to determine the degree to which risks from threatening factors could impact an entire population, and is not intended to be an estimate of the amount of occupied or potential habitat.
High threat exotic plant cover	Plant cover composed of vascular plants not native to Australia that if not controlled will invade and outcompete native plant species.
Hollow bearing tree	A living or dead tree that has at least one hollow. A tree is considered to contain a hollow if: (a) the entrance can be seen; (b) the minimum entrance width is at least 5 cm; (c) the hollow appears to have depth (i.e. you cannot see solid wood beyond the entrance); (d) the hollow is at least 1 m above the ground. Trees must be examined from all angles.
Important wetland	A wetland that is listed in the Directory of Important Wetlands of Australia (DIWA) and SEPP 14 Coastal Wetlands
Linear shaped development	Development that is generally narrow in width and extends across the landscape for a distance greater than 3.5 kilometres in length

Terminology	Definition
Local population	The population that occurs in the study area. In cases where multiple populations occur in the study area or a population occupies part of the study area, impacts on each subpopulation must be assessed separately.
Local wetland	Any wetland that is not identified as an important wetland (refer to definition of Important wetland).
NSW (Mitchell) landscape	Landscapes with relatively homogeneous geomorphology, soils and broad vegetation types, mapped at a scale of 1:250,000.
Multiple fragmentation impact development	Developments such as wind farms and coal seam gas extraction that require multiple extraction points (wells) or turbines and a network of associated development including roads, tracks, gathering systems/flow lines, transmission lines
Operational Manual	The Operational Manual published from time to time by DPIE, which is a guide to assist assessors when using the BAM
Patch size	An area of intact native vegetation that: a) occurs on the development site or biodiversity stewardship site, and b) includes native vegetation that has a gap of less than 100 m from the next area of native vegetation (or ≤ 30 m for non-woody ecosystems). Patch size may extend onto adjoining land that is not part of the development site or stewardship site..
Proponent	A person who intends to apply for consent to carry out development or for approval for an activity.
Reference sites	The relatively unmodified sites that are assessed to obtain local benchmark information when benchmarks in the Vegetation Benchmarks Database are too broad or otherwise incorrect for the PCT and/or local situation. Benchmarks can also be obtained from published sources.
Regeneration	The proportion of over-storey species characteristic of the PCT that are naturally regenerating and have a diameter at breast height < 5 cm within a vegetation zone.
Residual impact	An impact on biodiversity values after all reasonable measures have been taken to avoid, minimise or mitigate the impacts of development. Under the BAM, an offset requirement is determined for the remaining impacts on biodiversity values.
Retirement of credits	The purchase and retirement of biodiversity credits from an already-established biobank site or a biodiversity stewardship site secured by a biodiversity stewardship agreement.
Riparian buffer	Riparian buffers applied to water bodies in accordance with the BAM
Sensitive biodiversity values land map	Development within an area identified on the map requires assessment using the BAM.
Site attributes	The matters assessed to determine vegetation integrity. They include: native plant species richness, native over-storey cover, native mid-storey cover, native ground cover (grasses), native ground cover (shrubs), native ground cover (other), exotic plant cover (as a percentage of total ground and mid-storey cover), number of trees with hollows, proportion of over-storey species occurring as regeneration, and total length of fallen logs.
Site-based development	a development other than a linear shaped development, or a multiple fragmentation impact development
Species credits	The class of biodiversity credits created or required for the impact on threatened species that cannot be reliably predicted to use an area of land based on habitat surrogates. Species that require species credits are listed in the Threatened Biodiversity Data Collection.
Subject land	Is land to which the BAM is applied in Stage 1 to assess the biodiversity values of the land. It includes land that may be a development site, clearing site, proposed for biodiversity certification or land that is proposed for a biodiversity stewardship agreement.

Terminology	Definition
Threatened Biodiversity Data Collection	Part of the BioNet database, published by DPIE and accessible from the BioNet website.
Threatened species	Critically Endangered, Endangered or Vulnerable threatened species as defined by Schedule 1 of the BC Act, or any additional threatened species listed under Part 13 of the EPBC Act as Critically Endangered, Endangered or Vulnerable.
Vegetation Benchmarks Database	A database of benchmarks for vegetation classes and some PCTs. The Vegetation Benchmarks Database is published by DPIE and is part of the BioNet Vegetation Classification.
Vegetation zone	A relatively homogenous area of native vegetation on a development site, land to be biodiversity certified or a biodiversity stewardship site that is the same PCT and broad condition state.
Wetland	An area of land that is wet by surface water or ground water, or both, for long enough periods that the plants and animals in it are adapted to, and depend on, moist conditions for at least part of their life cycle. Wetlands may exhibit wet and dry phases and may be wet permanently, cyclically or intermittently with fresh, brackish or saline water
Woody native vegetation	Native vegetation that contains an over-storey and/or mid-storey that predominantly consists of trees and/or shrubs

Appendix B Vegetation Floristic Plot Data

Family	Species	Common Name	Listing Status	ROTAP	Exotic	High Threat Weed	Growth Form Group	Plot 1		
								Stratum & Layer	Cover	Abundance
Apiaceae	<i>Cyclospermum</i>	Slender			*			g	0.1	10
	<i>leptophyllum</i>	Celery								
Asteraceae	<i>Conyza</i>	Flaxleaf			*			g	0.1	10
	<i>bonariensis</i>	Fleabane								
Asteraceae	<i>Gamochaeta</i> spp.				*			g	0.5	100
Asteraceae	<i>Hypochaeris</i> <i>radicata</i>	Catsear			*			g	0.1	10
Asteraceae	<i>Senecio</i> <i>madagascariensis</i>	Fireweed			*	Yes		g	0.1	1
Asteraceae	<i>Soliva sessilis</i>	Bindyi			*			g	0.5	1000
Asteraceae	<i>Sonchus</i> <i>oleraceus</i>	Common Sowthistle			*			g	0.1	20
Asteraceae	<i>Taraxacum</i> <i>officinale</i>	Dandelion			*			g	0.1	10
Caryophyllaceae	<i>Cerastium</i> <i>glomeratum</i>	Mouse-ear Chickweed			*			g	0.2	20
Caryophyllaceae	<i>Petrorhagia</i> <i>dubia</i>				*			g	0.1	20
Caryophyllaceae	<i>Stellaria media</i>	Common Chickweed			*			g	0.2	500
Convolvulaceae	<i>Dichondra</i> <i>repens</i>	Kidney Weed					Forb (FG)	g	0.2	50
Fabaceae (Faboideae)	<i>Medicago</i> <i>sativa</i>	Lucerne			*			g	0.5	20
Fabaceae (Faboideae)	<i>Pultenaea</i> <i>microphylla</i>	A Bush Pea					Shrub (SG)	g	0.1	1
Malaceae	<i>Photinia</i> spp.				*			m	0.1	1
Myrtaceae	<i>Eucalyptus</i> <i>moluccana</i>	Grey Box					Tree (TG)	u	20	4
Oxalidaceae	<i>Oxalis</i> <i>perennans</i>						Forb (FG)	g	0.1	20
Plantaginaceae	<i>Plantago</i> <i>lanceolata</i>	Lamb's Tongues			*			g	0.5	50

Family	Species	Common Name	Listing Status	ROTAP	Exotic	High Threat Weed	Growth Form Group	Plot 1		
								Stratum & Layer	Cover	Abundance
Plantaginaceae	<i>Plantago myosuroides</i> subsp. <i>myosuroides</i>				*			g	0.5	100
Poaceae	<i>Briza subaristata</i>				*	Yes		g	0.1	20
Poaceae	<i>Bromus catharticus</i>	Prairie Grass			*			g	0.2	20
Poaceae	<i>Chloris truncata</i>	Windmill Grass					Grass & grasslike (GG)	g	0.1	10
Poaceae	<i>Cynodon dactylon</i>	Common Couch					Grass & grasslike (GG)	g	0.5	100
Poaceae	<i>Eragrostis curvula</i>	African Lovegrass			*	Yes		g	70	2000
Poaceae	<i>Paspalum dilatatum</i>	Paspalum			*	Yes		g	1	100
Poaceae	<i>Vulpia</i> spp.	Rat's-tail Fescue			*			g	10	500
Rubiaceae	<i>Richardia stellaris</i>				*			g	0.1	10

Appendix C Vegetation Integrity Plot Data

Plot location

Plot	PCT	Condition	Zone	Easting	Northing	Bearing
1	849	Low	56	295875	6257133	195°

Composition data

Tree	Shrub	Grass	Forbs	Ferns	Other
1	1	2	2	0	0

Structure data

Tree	Shrub	Grass	Forbs	Ferns	Other
20.0	0.1	0.6	0.3	0.0	0.0

Function data

Large Trees	Hollow trees	Litter Cover	Length of Fallen Logs	Tree Stem 5 to 9	Tree Stem 10 to 19	Tree Stem 20 to 29	Tree Stem 30 to 49	Tree Stem 50 to 79	Tree Stem 80+	Tree Regen	High Threat Exotic
0	0	16	0	0	0	1	1	0	0	0	71.2

Appendix D Biodiversity credit report



BAM Biodiversity Credit Report (Like for like)

Proposal Details

Assessment Id 00029977/BAAS18141/21/00029978	Proposal Name Chameleon Drive Erskine Park	BAM data last updated * 24/11/2021
Assessor Name Karen Leigh Spicer	Assessor Number BAAS18141	BAM Data version * 50
Proponent Names Fiona Kulak	Report Created 31/01/2022	BAM Case Status Finalised
Assessment Revision 0	Assessment Type Part 4 Developments (Small Area)	Date Finalised 31/01/2022
BOS entry trigger BOS Threshold: Biodiversity Values Map	* Disclaimer: BAM data last updated may indicate either complete or partial update of the BAM calculator database. BAM calculator database may not be completely aligned with Bionet.	

Potential Serious and Irreversible Impacts

Name of threatened ecological community	Listing status	Name of Plant Community Type/ID
Cumberland Plain Woodland in the Sydney Basin Bioregion	Critically Endangered Ecological Community	849-Cumberland shale plains woodland
Species		
Nil		

Additional Information for Approval

Assessment Id 00029977/BAAS18141/21/00029978	Proposal Name Chameleon Drive Erskine Park	Page 1 of 4
---	---	-------------



BAM Biodiversity Credit Report (Like for like)

PCT Outside Ibra Added
None added

PCTs With Customized Benchmarks

PCT No Changes

Predicted Threatened Species Not On Site

Name Grantiella picta / Painted Honeyeater

Ecosystem Credit Summary (Number and class of biodiversity credits to be retired)

Name of Plant Community Type/ID	Name of threatened ecological community	Area of impact	HBT Cr	No HBT Cr	Total credits to be retired
849-Cumberland shale plains woodland	Cumberland Plain Woodland in the Sydney Basin Bioregion	0.0	0	0	0

Assessment Id 00029977/BAAS18141/21/00029978	Proposal Name Chameleon Drive Erskine Park	Page 2 of 4
---	---	-------------



BAM Biodiversity Credit Report (Like for like)

849-Cumberland shale plains woodland	Like-for-like credit retirement options					
	Name of offset trading group	Trading group	Zone	HBT	Credits	IBRA region
	Cumberland Plain Woodland in the Sydney Basin Bioregion This includes PCT's: 849, 850	-	849_low	No	0	Cumberland, Burragorang, Pittwater, Sydney Cataract, Wollemi and Yengo. or Any IBRA subregion that is within 100 kilometers of the outer edge of the impacted site.

Species Credit Summary
No Species Credit Data

Credit Retirement Options Like-for-like credit retirement options

Assessment Id
00029977/BAAS18141/21/00029978

Proposal Name
Chameleon Drive Erskine Park

Page 3 of 4



BAM Biodiversity Credit Report (Like for like)

Assessment Id
00029977/BAAS18141/21/00029978

Proposal Name
Chameleon Drive Erskine Park

Page 4 of 4

Appendix E Likelihood of occurrence

An assessment of likelihood of occurrence was made for threatened and migratory species identified from the database search. Five terms for the likelihood of occurrence of species are used in this report. This assessment was based on database or other records, presence or absence of suitable habitat, features of the proposal site, results of the site inspection and professional judgement. Some Migratory or Marine species identified from the Commonwealth database search have been excluded from the assessment, due to lack of habitat. The terms for likelihood of occurrence are defined below:

- 'known' = the species was or has been observed on the site
- 'likely' = a medium to high probability that a species uses the site
- 'potential' = suitable habitat for a species occurs on the site, but there is insufficient information to categorise the species as likely to occur, or unlikely to occur
- 'unlikely' = a very low to low probability that a species uses the site
- 'no' = habitat within the subject site and in the vicinity is unsuitable for the species

Information provided in the habitat associations' column has primarily been extracted (and modified) from the Commonwealth Species Profile and Threats Database and the NSW Threatened Species Profiles.

Scientific Name	BC Act Status	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
ECOLOGICAL COMMUNITIES				
Coastal Swamp Oak (Casuarina glauca) Forest of New South Wales and South East Queensland ecological community		E	The ecological community occurs in sub-tropical, sub-humid and temperate climatic zones from Curtis Island, north of Gladstone, in Queensland to Bermagui in southern New South Wales. The ecological community is found within the South Eastern Queensland (SEQ), NSW North Coast (NNC), Sydney Basin (SYB) and South East Corner (SEC) IBRA7 Bioregions. Coastal Swamp Oak Forest is often found in association with other vegetation types such as coastal saltmarsh, mangroves, freshwater wetlands, littoral rainforests or swamp sclerophyll forests in a 'mosaic' of coastal floodplain communities.	No – this community was not identified within the subject site.
River-Flat Eucalypt Forest on Coastal Floodplains of the New South Wales North Coast, Sydney Basin and South East Corner Bioregions		CE	The structure of the community may vary from tall open forests (>40m) to woodlands. The most widespread and abundant dominant trees include Eucalyptus tereticornis (forest red gum), <i>E. amplifolia</i> (cabbage gum), <i>Angophora floribunda</i> (rough-barked apple) and <i>A. subvelutina</i> (broad-leaved apple). <i>Eucalyptus baueriana</i> (blue box), <i>E. botryoides</i> (bangalay) and <i>E. elata</i> (river peppermint) may be common south from Sydney. <i>E. ovata</i> (swamp gum) occurs on the far south coast, <i>E. saligna</i> (Sydney blue gum) and <i>E. grandis</i> (flooded gum) may occur north of Sydney, while <i>E. benthamii</i> is restricted to the Hawkesbury floodplain. A layer of small trees may be present, including <i>Melaleuca decora</i> , <i>M. styphelioides</i> (prickly-leaved teatree), <i>Backhousia myrtifolia</i> (grey myrtle), <i>Melia azadarach</i> (white cedar), <i>Casuarina cunninghamiana</i> (river oak) and <i>C. glauca</i> (swamp oak). Scattered shrubs include <i>Bursaria spinosa</i> , <i>Solanum prinophyllum</i> , <i>Rubus parvifolius</i> , <i>Breynia oblongifolia</i> , <i>Ozothamnus diosmifolius</i> , <i>Hymenanthera dentata</i> , <i>Acacia floribunda</i> and <i>Phyllanthus gunnii</i> . The groundcover is composed of abundant forbs, scramblers and grasses. Found on the river flats of the coastal floodplains. Known from parts of the Local Government Areas of Port Stephens, Maitland, Singleton, Cessnock, Lake Macquarie, Wyong, Gosford, Hawkesbury, Baulkham Hills, Blacktown, Parramatta, Penrith, Blue Mountains, Fairfield, Holroyd, Liverpool, Bankstown, Wollondilly, Camden, Campbelltown, Sutherland, Wollongong, Shellharbour, Kiama, Shoalhaven, Palerang, Eurobodalla and Bega Valley.	No – this community was not identified within the subject site.
Castlereagh Scribbly Gum and Agnes Banks Woodlands of the Sydney Basin Bioregion		E	Typically a low woodland, with canopy species reaching an average 15 m in height. The canopy is often dominated by one or more of <i>Angophora bakeri</i> (narrow leaved apple), <i>Eucalyptus racemosa</i> (narrow-leaved scribbly gum) and <i>E. parramattensis</i> subsp. <i>parramattensis</i> (Parramatta red gum). <i>Melaleuca</i> species including <i>M. decora</i> (paperbark)	No – this community was not identified within the subject site.

Scientific Name	BC Status	Act	EPBC Status	Act	Distribution and Habitat	Likelihood of Occurrence
					may also be prominent in the canopy and/or mid layer. The understorey has a prominent and diverse mid-layer of sclerophyll shrubs, sometimes dominated by either Banksia or Melaleuca species. It typically has a patchy ground cover of sedges and grasses. The isolation of the alluvial deposits in the Hawkesbury-Nepean river valley and differences in the soil characteristics have led to the development of differences in species composition and abundance across the range of the ecological community. In addition, the Agnes Banks vegetation occurs on aeolian sand and can contain a number of species reminiscent of communities closer to the coast, such as <i>Dillwynia glaberrima</i> , <i>Ricinocarpos pinifolius</i> (wedding bush) and <i>Banksia aemula</i> (wallum). Distributed throughout the Sydney Basin Bioregion, mostly in the Cumberland IBRA sub-region, with small occurrences in the Sydney Cataract, Wollemi and Burragorang sub-regions. It occurs primarily in the Castlereagh area in the north-west of the Cumberland Plain with other known occurrences near Holsworthy, Kemps Creek and Longneck Lagoon.	
Cumberland Plain Shale Woodlands and Shale-Gravel Transition Forest			CE		The minimum projected foliage cover of canopy trees is 10% or more and the tree canopy is typically dominated by <i>Eucalyptus moluccana</i> (Grey Box), <i>E. tereticornis</i> (Forest Red Gum) and/or <i>E. fibrosa</i> (Red Ironbark). A sparse lower tree layer may be present, typically with young eucalypts of upper tree canopy species and species of <i>Acacia</i> , <i>Exocarpos</i> and <i>Melaleuca</i> . The understorey typically is dominated by the ground layer, typically comprising a variety of perennial native graminoids and forbs. Endemic to the shale hills and plains of the Sydney Basin Bioregion in NSW, occurring primarily in, but not limited to, the Cumberland Sub-region.	No – this community was not identified within the subject site.
Cooks River/Castlereagh Ironbark Forest of the Sydney Basin Bioregion			CE		Ranges from open forest to low woodland, with a canopy dominated by <i>Eucalyptus fibrosa</i> (Broad-leaved Ironbark) and <i>Melaleuca decora</i> (Paperbark). The canopy may also include other eucalypts such as <i>E. longifolia</i> (Woollybutt). The dense shrubby understorey consists of <i>Melaleuca nodosa</i> (Prickly-leaved Paperbark) and <i>Lissanthe strigosa</i> (Peach Heath), with a range of 'pea' flower shrubs, such as <i>Dillwynia tenuifolia</i> , <i>Pultenaea villosa</i> (Hairy Bush-pea) and <i>Daviesia ulicifolia</i> (Gorse Bitter Pea). The sparse ground layer contains a range of grasses and herbs. Occurs in western Sydney, with the most extensive stands occurring in the Castlereagh and Holsworthy areas. Smaller remnants occur in the Kemps Creek area and in the eastern section of the Cumberland Plain.	No – this community was not identified within the subject site.

Scientific Name	BC Status	Act	EPBC Status	Act	Distribution and Habitat	Likelihood of Occurrence
Western Sydney Dry Rainforest and Moist Woodland on Shale			CE		Typically a low closed forest, slightly more open in the moist woodland form, with emergent trees up to 25 m high and a lower tree layer. In sheltered gullies and on lower slopes the canopy layer is typically dominated by <i>Melaleuca styphelioides</i> (prickly-leaved paperbark). Other diagnostic tree species include <i>Acacia implexa</i> (hickory wattle), <i>Alectryon subcinereus</i> (native quince), <i>Brachychiton populneus</i> (kurrajong), <i>Corymbia maculata</i> (spotted gum), <i>Melicope micrococca</i> (white euodia) and <i>Streblus pendulinus</i> (whalebone tree). Generally on upper slopes to undulating terrain, or at more disturbed sites, the ecological community exhibits its moist woodland form with the canopy dominated by <i>E. moluccana</i> , <i>E. tereticornis</i> , <i>E. crebra</i> and/or <i>Corymbia maculata</i> . Characteristic shrub species include <i>Breynia oblongifolia</i> (false coffee bush), <i>Clerodendrum tomentosum</i> (hairy clerodendrum) and <i>Notelaea longifolia</i> f. <i>longifolia</i> (large mock-olive). Vines and other climber species are typically common. The ground layer is variable and generally sparse with a diverse mix of forbs, ferns and shade-tolerant grasses. Distributed throughout the Cumberland Plain Sub-region of the Sydney Basin Bioregion.	No – this community was not identified within the subject site.

Scientific Name	Common Name	BC Status	Act	EPBC Status	Act	Distribution and Habitat	Likelihood of Occurrence
FAUNA							
<i>Actitis hypoleucos</i>	Common Sandpiper			M		Summer migrant. In NSW, widespread along coastline and also occurs in many areas inland. Coastal wetlands and some inland wetlands, especially muddy margins or rocky shores. Also estuaries and deltas, lakes, pools, billabongs, reservoirs, dams and claypans, mangroves.	No – no habitat within the subject site.
<i>Anthochaera phrygia</i>	Regent Honeyeater	CE		CE		Inland slopes of south-east Australia, and less frequently in coastal areas. In NSW, most records are from the North-West Plains, North-West and South-West Slopes, Northern Tablelands, Central Tablelands and Southern Tablelands regions; also recorded in the Central Coast and Hunter Valley regions. Eucalypt woodland and open forest, wooded farmland and urban areas with mature eucalypts, and riparian forests of <i>Casuarina cunninghamiana</i> (River Oak).	Unlikely – lack of suitable habitat present within the subject site.

Scientific Name	Common Name	BC Status	Act	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
<i>Apus pacificus</i>	Fork-tailed Swift			M	Recorded in all regions of NSW. Riparian woodland., swamps, low scrub, heathland, saltmarsh, grassland, Spinifex sandplains, open farmland and inland and coastal sand-dunes.	Unlikely – lack of suitable habitat present within the subject site.
<i>Botaurus poiciloptilus</i>	Australasian Bittern	E		E	Found over most of NSW except for the far north-west. Permanent freshwater wetlands with tall, dense vegetation, particularly Typha spp. (bullrushes) and Eleocharis spp. (spikerushes).	No – no suitable habitat within the subject site.
<i>Calidris acuminata</i>	Sharp-tailed Sandpiper			M	Summer migrant. Widespread in most regions of NSW, especially in coastal areas, but sparse in the south-central Western Plain and east Lower Western Regions. Shallow fresh or brackish wetlands, with inundated or emergent sedges, grass, saltmarsh or other low vegetation.	No – no suitable habitat within the subject site.
<i>Calidris melanotos</i>	Pectoral Sandpiper			M	Summer migrant to Australia. Widespread but scattered in NSW. East of the Great Divide, recorded from Casino and Ballina, south to Ulladulla. West of the Great Divide, widespread in the Riverina and Lower Western regions. Shallow fresh to saline wetlands, including coastal lagoons, estuaries, bays, swamps, lakes, inundated grasslands, saltmarshes, river pools, creeks, floodplains and artificial wetlands.	No – no suitable habitat within the subject site.
<i>Chalinolobus dwyeri</i>	Large-eared Pied Bat	V		V	Recorded from Rockhampton in Qld south to Ulladulla in NSW. Largest concentrations of populations occur in the sandstone escarpments of the Sydney basin and the NSW north-west slopes. Wet and dry sclerophyll forests, Cyprus Pine dominated forest, woodland, sub-alpine woodland, edges of rainforests and sandstone outcrop country.	Unlikely – lack of breeding habitat within the subject site. Degraded foraging habitat unlikely to support this species.
<i>Dasyurus maculatus maculatus</i> (SE mainland population)	Spotted-tailed Quoll	V		E	Found on the east coast of NSW, Tasmania, eastern Victoria and north-eastern Qld. Rainforest, open forest, woodland, coastal heath and inland riparian forest, from the sub-alpine zone to the coastline.	Unlikely – lack of suitable habitat present within the subject site.
<i>Falco hypoleucos</i>	Grey Falcon	E1			Arid and semi-arid zones. In NSW, found chiefly throughout the Murray-Darling Basin, with the occasional vagrant east of the Great Dividing Range. Shrubland, grassland and	Unlikely – lack of suitable habitat

Scientific Name	Common Name	BC Status	Act	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
					wooded watercourses, occasionally in open woodlands near the coast, and near wetlands.	present within the subject site.
<i>Gallinago hardwickii</i>	Latham's Snipe			M	Migrant to east coast of Australia, extending inland west of the Great Dividing Range in NSW. Freshwater, saline or brackish wetlands up to 2000 m above sea-level; usually freshwater swamps, flooded grasslands or heathlands.	Unlikely – lack of suitable habitat present within the subject site.
<i>Grantiella picta</i>	Painted Honeyeater	V		V	Widely distributed in NSW, predominantly on the inland side of the Great Dividing Range but avoiding arid areas. Boree, Brigalow and Box-Gum Woodlands and Box-Ironbark Forests.	No – lack of suitable habitat present within the subject site.
<i>Heleioporus australiacus</i>	Giant Burrowing Frog	V		V	South eastern NSW and Victoria, in two distinct populations: a northern population in the sandstone geology of the Sydney Basin as far south as Ulladulla, and a southern population occurring from north of Narooma through to Walhalla, Victoria. Heath, woodland and open dry sclerophyll forest on a variety of soil types except those that are clay based.	No – lack of suitable habitat present within the subject site.
<i>Lathamus discolor</i>	Swift Parrot	E		CE	Migrates from Tasmania to mainland in Autumn-Winter. In NSW, the species mostly occurs on the coast and south west slopes. Box-ironbark forests and woodlands.	No – lack of suitable habitat present within the subject site.
<i>Litoria aurea</i>	Green and Golden Bell Frog	E		V	Since 1990, recorded from ~50 scattered sites within its former range in NSW, from the north coast near Brunswick Heads, south along the coast to Victoria. Records exist west to Bathurst, Tumut and the ACT region. Marshes, dams and stream-sides, particularly those containing Typha spp. (bullrushes) or Eleocharis spp. (spikerushes). Some populations occur in highly disturbed areas.	Unlikely – lack of suitable habitat present within the subject site.
<i>Monarcha melanopsis</i>	Black-faced Monarch			M	In NSW, occurs around the eastern slopes and tablelands of the Great Divide, inland to Coutts Crossing, Armidale, Widden Valley, Wollemi National Park and Wombeyan Caves. It is rarely recorded farther inland. Rainforest, open eucalypt forests, dry sclerophyll forests and woodlands, gullies in mountain areas or coastal foothills, Brigalow scrub, coastal scrub, mangroves, parks and gardens.	Unlikely – lack of suitable habitat present within the subject site.

Scientific Name	Common Name	BC Act Status	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
<i>Motacilla flava</i>	Yellow Wagtail		M	Regular summer migrant to mostly coastal Australia. In NSW recorded Sydney to Newcastle, the Hawkesbury and inland in the Bogan LGA. Swamp margins, sewage ponds, saltmarshes, playing fields, airfields, ploughed land, lawns.	Unlikely – lack of suitable habitat present within the subject site.
<i>Myiagra cyanoleuca</i>	Satin Flycatcher		M	In NSW, widespread on and east of the Great Divide and sparsely scattered on the western slopes, with very occasional records on the western plains. Eucalypt-dominated forests, especially near wetlands, watercourses, and heavily-vegetated gullies.	Unlikely – lack of suitable habitat present within the subject site.
<i>Pandion haliaetus</i>	Osprey		M	The breeding range of the Eastern Osprey extends around the northern coast of Australia (including many offshore islands) from Albany in Western Australia to Lake Macquarie in NSW. Eastern Ospreys occur in littoral and coastal habitats and terrestrial wetlands of tropical and temperate Australia and offshore islands. They are mostly found in coastal areas but occasionally travel inland along major rivers, particularly in northern Australia. They require extensive areas of open fresh, brackish or saline water for foraging. They frequent a variety of wetland habitats including inshore waters, reefs, bays, coastal cliffs, beaches, estuaries, mangrove swamps, broad rivers, reservoirs and large lakes and waterholes. They exhibit a preference for coastal cliffs and elevated islands in some parts of their range, but may also occur on low sandy, muddy or rocky shores and over coral cays. They may occur over atypical habitats such as heath, woodland or forest when travelling to and from foraging sites.	Unlikely – lack of suitable habitat present within the subject site.
<i>Petauroides volans</i>	Greater Glider population in the Eurobodalla local government area		V	This population on the south coast of NSW is bounded by the Moruya River to the north, Coila Lake to the south and the Princes Highway and cleared land exceeding 700 m in width to the west. Eucalypt forests and woodlands.	Unlikely – lack of suitable habitat present within the subject site.
<i>Petrogale penicillata</i>	Brush-tailed Rock-wallaby	E	V	In NSW they occur from the Qld border in the north to the Shoalhaven in the south, with the population in the Warrumbungle Ranges being the western limit. Rocky escarpments, outcrops and cliffs with a preference for complex structures with fissures, caves and ledges.	No – lack of suitable habitat present within the subject site.

Scientific Name	Common Name	BC Act Status	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
<i>Phascolarctos cinereus</i>	Koala, Hawks Nest and Tea Gardens population	E,V	V	Known from, and in the immediate vicinity of, the towns of Hawks Nest and Tea Gardens in the Great Lakes Local Government Area. Eucalypt forest and woodland communities, including coastal forests, rainforest, riparian areas, swamp sclerophyll forests, heathland and shrubland.	No – outside of natural range and lack of suitable habitat present within the subject site.
<i>Phascolarctos cinereus</i>	Koala in the Pittwater Local Government Area	E, V	V	The endangered population occurs within the Pittwater Local Government Area, with most recent records occurring on the Barrenjoey Peninsula. Eucalypt forests and woodlands. Key likely habitats within Pittwater Council are: Swamp Mahogany Forest, ecotone between Spotted Gum Forest & Hawkesbury Sandstone Open-Forest, Northern form of Coastal Sandstone Woodland at Whale Beach, Red Bloodwood - Scribbly Gum Woodland, Bilgola Plateau Forest and the Grey Ironbark - Grey Gum form of the Newport Bangalay Woodland.	No – outside of natural range and lack of suitable habitat present within the subject site.
<i>Phascolarctos cinereus</i>	Koala	V	V	In NSW it mainly occurs on the central and north coasts with some populations in the west of the Great Dividing Range. There are sparse and possibly disjunct populations in the Bega District, and at several sites on the southern tablelands. Eucalypt woodlands and forests.	Unlikely – no individuals were observed during the site visit and there is a lack of suitable habitat present within the subject site. Lack of connectivity to known habitat of this species.
<i>Pseudomys novaehollandiae</i>	New Holland Mouse		V	Fragmented distribution across eastern NSW. Open heathlands, woodlands and forests with a heathland understorey, vegetated sand dunes.	No – lack of suitable habitat present within the subject site.
<i>Pteropus poliocephalus</i>	Grey-headed Flying-fox	V	V	Along the eastern coast of Australia, from Bundaberg in Qld to Melbourne in Victoria. Subtropical and temperate rainforests, tall sclerophyll forests and woodlands, heaths and swamps as well as urban gardens and cultivated fruit crops.	Potential – marginal foraging habitat

Scientific Name	Common Name	BC Status	Act	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
						within the subject site.
<i>Rhipidura rufifrons</i>	Rufous Fantail			M	Coastal and near coastal districts of northern and eastern Australia, including on and east of the Great Divide in NSW. Wet sclerophyll forests, subtropical and temperate rainforests. Sometimes drier sclerophyll forests and woodlands.	Unlikely – lack of suitable habitat present within the subject site.
<i>Rostratula australis</i>	Australian Painted Snipe	E		E	In NSW most records are from the Murray-Darling Basin. Other recent records include wetlands on the Hawkesbury River and the Clarence and lower Hunter Valleys. Swamps, dams and nearby marshy areas.	Unlikely – lack of suitable habitat present within the subject site.
<i>Tringa nebularia</i>	Common Greenshank			M	Summer migrant to Australia. Recorded in most coastal regions of NSW; also widespread west of the Great Dividing Range, especially between the Lachlan and Murray Rivers and the Darling River drainage basin, including the Macquarie Marshes, and north-west regions. Terrestrial wetlands (swamps, lakes, dams, rivers, creeks, billabongs, waterholes and inundated floodplains, claypans, saltflats, sewage farms and saltworks dams, inundated rice crops and bores) and sheltered coastal habitats (mudflats, saltmarsh, mangroves, embayments, harbours, river estuaries, deltas, lagoons, tidal pools, rock-flats and rock platforms).	No – lack of suitable habitat present within the subject site.
FLORA						
<i>Acacia bynoeana</i>	Bynoe's Wattle	E		V	Found in central eastern NSW, from the Hunter District (Morisset) south to the Southern Highlands and west to the Blue Mountains. Heath or dry sclerophyll forest on sandy soils.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Acacia pubescens</i>	Downy Wattle	V		V	Restricted to the Sydney region around the Bankstown-Fairfield-Rookwood and Pitt Town area, with outliers occurring at Barden Ridge, Oakdale and Mountain Lagoon. Open woodland and forest, including Cooks River/Castlereagh Ironbark Forest, Shale/Gravel Transition Forest and Cumberland Plain Woodland. Occurs on alluviums, shales and at the intergrade between shales and sandstones.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.

Scientific Name	Common Name	BC Status	Act	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
<i>Allocasuarina glareicola</i>		E		E	Primarily restricted to the Richmond (NW Cumberland Plain) district, but with an outlier population found at Voyager Point, Liverpool. Castlereagh woodland on lateritic soil. Found in open woodland with <i>Eucalyptus parramattensis</i> , <i>Eucalyptus fibrosa</i> , <i>Angophora bakeri</i> , <i>Eucalyptus sclerophylla</i> and <i>Melaleuca decora</i> .	No – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Cynanchum elegans</i>	White-flowered Wax Plant	E		E	Restricted to eastern NSW, from Brunswick Heads on the north coast to Gerroa in the Illawarra region, and as far west as Merriwa in the upper Hunter River valley. Dry rainforest; littoral rainforest; <i>Leptospermum laevigatum</i> - <i>Banksia integrifolia</i> subsp. <i>integrifolia</i> (Coastal Tea-tree– Coastal Banksia) coastal scrub; <i>Eucalyptus tereticornis</i> (Forest Red Gum) or <i>Corymbia maculata</i> (Spotted Gum) open forest and woodland; and <i>Melaleuca armillaris</i> (Bracelet Honey myrtle) scrub.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Genoplesium baueri</i>	Bauer's Midge Orchid	E		E	Has been recorded from locations between Nowra and Pittwater and may occur as far north as Port Stephens. Dry sclerophyll forest and moss gardens over sandstone.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Grevillea parviflora</i> subsp. <i>parviflora</i>	Small-flower Grevillea	V		V	Sporadically distributed throughout the Sydney Basin and in the Hunter in the Cessnock - Kurri Kurri area. Also known from Putty to Wyong and Lake Macquarie on the Central Coast. Heath and shrubby woodland to open forest on sandy or light clay soils usually over thin shales.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Haloragis exalata</i> subsp. <i>exalata</i>	Square Raspwort	V		V	Disjunct distribution in the Central Coast, South Coast and North Western Slopes botanical subdivisions of NSW. Protected and shaded damp situations in riparian habitats.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.

Scientific Name	Common Name	BC Status	Act	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
<i>Melaleuca deanei</i>	Deane's Paperbark	V		V	Ku-ring-gai/Berowra area, Holsworthy/Wedderburn area, Springwood (in the Blue Mountains), Wollemi National Park, Yalwal (west of Nowra) and Central Coast (Hawkesbury River) areas. Heath on sandstone.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Micromyrtus minutiflora</i>		E1		V	Restricted to the general area between Richmond and Penrith, Western Sydney. Castlereagh Scribbly Gum Woodland, Ironbark Forest, Shale/Gravel Transition Forest, open forest on tertiary alluvium and consolidated river sediments.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Persicaria elatior</i>	Tall Knotweed	V		V	In south-eastern NSW recorded from Mt Dromedary, Moruya State Forest near Turlinjah, the Upper Avon River catchment north of Robertson, Bermagui, and Picton Lakes. In northern NSW known from Raymond Terrace (near Newcastle) and the Grafton area (Cherry Tree and Gibberagee State Forests). Beside streams and lakes, swamp forest or disturbed areas.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Persoonia hirsuta</i>	Hairy Geebung	E		E	Scattered distribution around Sydney, from Singleton in the north, along the east coast to Bargo in the south and the Blue Mountains to the west. Sandy soils in dry sclerophyll open forest, woodland and heath on sandstone.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Persoonia nutans</i>	Nodding Geebung	E		E	Restricted to the Cumberland Plain in western Sydney, between Richmond in the north and Macquarie Fields in the south. Northern populations: sclerophyll forest and woodland (Agnes Banks Woodland, Castlereagh Scribbly Gum Woodland and Cooks River / Castlereagh Ironbark Forest) on aeolian and alluvial sediments. Southern populations: tertiary alluvium, shale sandstone transition communities and Cooks River / Castlereagh Ironbark Forest.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.

Scientific Name	Common Name	BC Status	Act	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
<i>Pimelea curviflora</i> var. <i>curviflora</i>		V		V	Confined to the coastal area of the Sydney and Illawarra regions between northern Sydney and Maroota in the north-west and Croom Reserve near Albion Park in the south. Woodland, mostly on shaley/lateritic soils over sandstone and shale/sandstone transition soils on ridgetops and upper slopes.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Pimelea spicata</i>	Spiked Rice-flower	E		E	Two disjunct areas; the Cumberland Plain (Marayong and Prospect Reservoir south to Narellan and Douglas Park) and the Illawarra (Landsdowne to Shellharbour to northern Kiama). Well-structured clay soils. Eucalyptus moluccana (Grey Box) communities and in areas of ironbark on the Cumberland Plain. Coast Banksia open woodland or coastal grassland in the Illawarra.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Pomaderris brunnea</i>	Brown Pomaderris	E		V	In NSW, found around the Colo, Nepean and Hawkesbury Rivers, including the Bargo area and near Camden. It also occurs near Walcha on the New England tablelands. Moist woodland or forest on clay and alluvial soils of flood plains and creek lines.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Pterostylis gibbosa</i>	Illawarra Greenhood	E		E	Known from a small number of populations in the Hunter region (Milbrodale), the Illawarra region (Albion Park and Yallah) and the Shoalhaven region (near Nowra). Open forest or woodland, on flat or gently sloping land with poor drainage.	Unlikely – the subject site is not within the known distribution of this species. The habitat on site is degraded. This species was not identified during the field survey.
<i>Pterostylis saxicola</i>	Sydney Plains Greenhood	E		E	Restricted to western Sydney between Freemans Reach in the north and Picton in the south. Small pockets of shallow soil in depressions on sandstone rock shelves above cliff lines, adjacent to sclerophyll forest or woodland on shale/sandstone transition soils or shale soils.	No – lack of suitable habitat present within the subject site. Not identified

Scientific Name	Common Name	BC Status	Act	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
						during the field survey.
<i>Pultenaea parviflora</i>		E		V	Endemic to the Cumberland Plain. Mainly from Windsor to Penrith and east to Dean Park, with outlier populations at Kemps Creek and Wilberforce. Dry sclerophyll forest, especially Castlereagh Ironbark Forest, Shale Gravel Transition Forest and transitional areas where these communities adjoin Castlereagh Scribbly Gum Woodland.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Rhizanthella slateri</i>	Rhizanthella slateri (Rupp) M.A. Clem. & Cribb in the Great Lakes local government area	E,V		E	The population occurs near Bulahdelah (within the Great Lakes LGA). Sclerophyll forest in shallow to deep loams.	No – outside of natural range and lack of suitable habitat present within the subject site.
<i>Rhizanthella slateri</i>	Eastern Australian Underground Orchid	V		E	In NSW, currently known from fewer than 10 locations, including near Bulahdelah, the Watagan Mountains, the Blue Mountains, Wiseman's Ferry area, Agnes Banks and near Nowra. Sclerophyll forest in shallow to deep loams.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Rhodamnia rubescens</i>	Scrub Turpentine	CE		CE	Occurs in coastal districts north from Batemans Bay in New South Wales, approximately 280 km south of Sydney, to areas inland of Bundaberg in Queensland. Populations of <i>R. rubescens</i> typically occur in coastal regions and occasionally extend inland onto escarpments up to 600 m a.s.l. in areas with rainfall of 1,000-1,600 mm. Found in littoral, warm temperate and subtropical rainforest and wet sclerophyll forest usually on volcanic and sedimentary soils.	No – lack of suitable habitat present within the subject site. Not identified during the field survey.
<i>Syzygium paniculatum</i>	Magenta Lilly Pilly	E		V	Only in NSW, in a narrow, linear coastal strip from Upper Lansdowne to Conjola State Forest. Subtropical and littoral rainforest on gravels, sands, silts and clays.	Unlikely – lack of suitable habitat present within the

Scientific Name	Common Name	BC Act Status	EPBC Act Status	Distribution and Habitat	Likelihood of Occurrence
					subject site. Not identified during the field survey.
<i>Thesium australe</i>	Austral Toadflax	V	V	In eastern NSW it is found in very small populations scattered along the coast, and from the Northern to Southern Tablelands. Grassland on coastal headlands or grassland and grassy woodland away from the coast.	Unlikely – lack of suitable habitat present within the subject site. Not identified during the field survey.

